

HÖLDER STABLY DETERMINING THE TIME-DEPENDENT ELECTROMAGNETIC POTENTIAL OF THE SCHRÖDINGER EQUATION

YAVAR KIAN* AND ERIC SOCCORSI*

ABSTRACT. We consider the inverse problem of determining the time- and space-dependent electromagnetic potential of the Schrödinger equation in a bounded domain of \mathbb{R}^n , $n \geq 2$, by boundary observation of the solution over the entire time span. In contrast with published mathematical results on the recovery of unknown coefficients depending on all variables of the Cauchy problem under consideration, which, to our best knowledge, are of logarithmic type, we establish upon assuming that the divergence of the magnetic potential is fixed, that the electric potential and the magnetic potential can be Hölder stably retrieved from those lateral measurements.

Keywords: Inverse problem, stability estimate, Schrödinger equation, time-dependent electromagnetic potential.

Mathematics subject classification 2010: 35R30, 35Q41.

1. INTRODUCTION

1.1. Statement of the problem. Let Ω be a bounded, simply connected domain of \mathbb{R}^n , $n \geq 2$, with C^3 boundary $\partial\Omega$. For $T \in (0, +\infty)$, we consider the initial boundary value problem (IBVP)

$$\begin{cases} (i\partial_t + \Delta_A + q)u = 0 & \text{in } Q := (0, T) \times \Omega, \\ u(0, \cdot) = 0 & \text{in } \Omega, \\ u = g & \text{on } \Sigma := (0, T) \times \Gamma, \end{cases} \quad (1.1)$$

where Δ_A is the Laplace operator $(\nabla + iA(t, x)) \cdot (\nabla + iA(t, x))$, associated with the real-valued magnetic potential $A := (a_j)_{1 \leq j \leq n} \in W^{1, \infty}(Q; \mathbb{R})^n$, i.e.

$$\Delta_A := \sum_{j=1}^n (\partial_{x_j} + ia_j(t, x))^2 = \Delta + 2iA(t, x) \cdot \nabla + i(\nabla \cdot A(t, x)) - |A(t, x)|^2 \quad (1.2)$$

and $q \in L^\infty(Q; \mathbb{R})$ is a real-valued electric potential. Here and in the remaining part of this text, we denote by $\nabla := (\partial_{x_1}, \dots, \partial_{x_n})^T$ the gradient operator with respect to the spatial variable $x := (x_1, \dots, x_n) \in \mathbb{R}^n$, the symbol \cdot (resp., $|\cdot|$) stands for the Euclidian scalar product (resp., norm) in \mathbb{R}^n , and the divergence operator with respect to $x \in \mathbb{R}^n$ is represented by the notation $\nabla \cdot$.

For all $s, r \in (0, +\infty)$ and for X being either Ω or $\partial\Omega$, we equip the functional spaces $H^{r,s}((0, T) \times X) := H^r(0, T; L^2(X)) \cap L^2(0, T; H^s(X))$ with the following norm

$$\|u\|_{H^{r,s}((0,T) \times X)}^2 := \|u\|_{H^r(0,T; L^2(X))}^2 + \|u\|_{L^2(0,T; H^s(X))}^2,$$

and we write $H^{r,s}(Q)$ (resp., $H^{r,s}(\Sigma)$) instead of $H^{r,s}((0, T) \times \Omega)$ (resp., $H^{r,s}((0, T) \times \partial\Omega)$). Then, for all

$$g \in \mathcal{H}(\Sigma) := \left\{ g \in H^{\frac{5}{2}, \frac{5}{2}}(\Sigma); g(0, \cdot) = \partial_t g(0, \cdot) = 0 \text{ on } \partial\Omega \right\},$$

* Aix Marseille Université, Université de Toulon, CNRS, CPT, Marseille, France.

we establish in Proposition 2.1 below, that there exists a unique solution $u_g \in H^{1,2}(Q)$ to (1.1) and that the mapping $g \mapsto u_g$ is continuous. As a corollary the Dirichlet-to-Neumann (DN) operator associated with (1.1), defined by

$$\begin{aligned} \Lambda_{A,q} : \mathcal{H}(\Sigma) &\rightarrow L^2(\Sigma) \\ g &\mapsto (\partial_\nu + iA \cdot \nu)u_g, \end{aligned} \tag{1.3}$$

where $\nu(x)$ denotes the unit outward normal to $\partial\Omega$ at x , is bounded. The main purpose of this paper is to examine the stability issue in the inverse problem of determining the electromagnetic potential (A, q) from the knowledge of $\Lambda_{A,q}$.

However, there is a natural obstruction to uniqueness in this problem, in the sense that the mapping $(A, q) \mapsto \Lambda_{A,q}$ is not injective. This can be seen from the identity

$$i\partial_t + \Delta_A + q = e^{-i\phi}(i\partial_t + \Delta_{\tilde{A}} + \tilde{q})e^{i\phi}, \quad \phi \in W^{3,\infty}(Q)$$

arising from (1.2) with $(\tilde{A}, \tilde{q}) = G_{A,q}(\phi) := (A - \nabla\phi, q + \partial_t\phi)$, which entails that $(i\partial_t + \Delta_{\tilde{A}} + \tilde{q})e^{i\phi}u_g = e^{i\phi}(i\partial_t + \Delta_A + q)u_g = 0$ for all $g \in \mathcal{H}(\Sigma)$. Therefore, if ϕ vanishes on Σ then it is apparent that $e^{i\phi}u_g$ is the $H^{1,2}(Q)$ -solution to (1.1), where (\tilde{A}, \tilde{q}) is substituted for (A, q) . Consequently, it holds true that

$$\Lambda_{\tilde{A},\tilde{q}}g = (\partial_\nu + i\tilde{A} \cdot \nu)e^{i\phi}u_g = e^{i\phi}(\partial_\nu + iA \cdot \nu)u_g = \Lambda_{A,q}g,$$

despite of the fact that (\tilde{A}, \tilde{q}) does not coincide with (A, q) whenever ϕ is not uniformly zero in Q . Otherwise stated, since the DN map (1.3) is invariant under the gauge transformation $(A, q) \mapsto G_{A,q}(\phi)$ associated with $\phi \in W_*^{3,\infty}(Q) := \{\phi \in W^{3,\infty}(Q); \phi|_\Sigma = 0\}$, then it is hopeless to retrieve (A, q) through $\Lambda_{A,q}$ and the best we can expect is to determine the *gauge class* $G_{A,q}(W_*^3(Q)) := \{G_{A,q}(\phi), \phi \in W_*^{3,\infty}(Q)\}$ of (A, q) . Moreover, for any two gauge equivalent electromagnetic potentials (A, q) and (\tilde{A}, \tilde{q}) , there exists a unique $\phi \in W_*^{3,\infty}(Q)$ such that we have $(\tilde{A}, \tilde{q}) = G_{A,q}(\phi)$ and we notice for each for $t \in (0, T)$ that the function $\phi(t, \cdot)$ is solution to the following elliptic system:

$$\begin{cases} -\Delta\phi(t, \cdot) = \nabla \cdot (\tilde{A} - A)(t, \cdot) & \text{in } \Omega, \\ \phi(t, \cdot) = 0 & \text{on } \partial\Omega. \end{cases}$$

Therefore, if the time-dependent electromagnetic potential (A, q) can be determined modulo gauge invariance by $\Lambda_{A,q}$ then it is actually possible to recover (A, q) itself provided the divergence $\nabla \cdot A$ is known.

1.2. What is known so far. Since inverse problems are of great interest in applied sciences, it is no surprise that the determination of coefficients in partial differential equations such as the magnetic Schrödinger equation under study in this article has attracted the attention of numerous mathematicians over the previous decades.

For instance, using the Bukhgeim-Klibanov method [15], Baudouin and Puel [3] proved Lipschitz stable identification of the time independent electric potential in the dynamical (i.e., non stationary) Schrödinger equation from a single boundary observation of the solution. Here the measurement can be performed on any subpart of the boundary fulfilling the geometric control property expressed by Bardos, Lebeau and Rauch in [2]. This condition was removed by Bellassoued and Choulli in [6], provided the electric potential is *a priori* known in a neighborhood of the boundary. We refer to [19] for the Lipschitz stable reconstruction of the magnetic potential in the Coulomb gauge class by a finite number of boundary measurements of the solution to the Schrödinger equation. More recently, in [13], Ben Aïcha and Mejri claimed simultaneous Lipschitz stable determination of the electric potential and the divergence free magnetic potential, from the same type of boundary data.

The above mentioned results involve a finite number of boundary observations of the solution, performed over the entire time span. But this is no longer the case in [7] where the magnetic field was stably recovered from the knowledge of the DN map associated with the dynamic Schrödinger equation. In the same spirit, Bellassoued and Dos Santos Ferreira proved stable identification of the electric potential by the DN map associated with the Schrödinger equation on a Riemannian manifold in [8]. This result was extended in [4] to simultaneous determination of the electric potential and the magnetic field. We also refer to [20, 24, 32] for an extensive treatment of similar inverse problems. We stress out that all the above results were established in a bounded domain and that the analysis carried out in [3] (resp. [6] and [7]) was adapted to the case of unbounded cylindrical domains in [9] (resp., [10] and [35, 36]).

All the above mentioned works are concerned with space-only dependent (i.e. time independent) coefficients. Actually, there is only a very small number of papers available in the mathematical literature, dealing with the inverse problem of determining time-dependent coefficients of the Schrödinger equation. For instance, it was proved in [21] that the DN map uniquely determines the time-dependent electromagnetic potential modulo gauge invariance. The stability issue for the same problem was examined in [18], where the time-dependent electric potential was logarithmic stably recovered from boundary observation for all times and internal measurement at final time, of the solution. More recently, in [12], this approach was adapted to the case of an electromagnetic potential with sufficiently small time independent magnetic part. To the best of our best knowledge, these two last articles are the only mathematical papers studying the stability issue in the inverse problem of determining time-dependent coefficients of the Schrödinger equation. Nevertheless, we point out that similar problems were addressed in [5, 11, 16, 17, 22, 23, 26, 27, 28, 30, 31, 33, 41, 40, 42] for either parabolic, hyperbolic, or even time-fractional diffusion equations.

1.3. Main result. The main result of this paper is the following Hölder stability estimate of the electromagnetic potential entering the Schrödinger equation in (1.1), with respect to the DN map.

Theorem 1.1. *Fix $M \in (0, +\infty)$ and for $j = 1, 2$, let $A_j \in W^{5,\infty}(Q)^n \cap H^6(Q)^n$ and $q_j \in W^{4,\infty}(Q)$ satisfy the three following conditions:*

$$\partial_x^\alpha A_1(t, x) = \partial_x^\alpha A_2(t, x), \quad (t, x) \in \Sigma, \quad \alpha \in \mathbb{N}^n, \quad |\alpha| \leq 5, \quad (1.4)$$

$$\nabla \cdot A_1(t, x) = \nabla \cdot A_2(t, x), \quad (t, x) \in Q \quad (1.5)$$

and

$$\sum_{j=1}^2 (\|A_j\|_{W^{5,\infty}(Q)^n} + \|A_j\|_{H^6(Q)^n} + \|q_j\|_{W^{4,\infty}(Q)}) \leq M. \quad (1.6)$$

Then, there exist three positive constants, r and s , depending only on n , and C , depending only on n, T, Ω and M , such that we have

$$\|A_1 - A_2\|_{L^2(0,T;H^5(\Omega))} \leq C \|\Lambda_{A_1, q_1} - \Lambda_{A_2, q_2}\|^r \quad (1.7)$$

and

$$\|q_1 - q_2\|_{H^{-1}(Q)} \leq C \|\Lambda_{A_1, q_1} - \Lambda_{A_2, q_2}\|^s. \quad (1.8)$$

In Theorem 1.1 and the remaining part of this article, the DN maps Λ_{A_j, q_j} , $j = 1, 2$, lie in the space $\mathcal{B}(\mathcal{H}(\Sigma), L^2(\Sigma))$ of linear bounded operators from $\mathcal{H}(\Sigma)$ into $L^2(\Sigma)$ and $\|\cdot\|$ denotes the usual norm in $\mathcal{B}(\mathcal{H}(\Sigma), L^2(\Sigma))$.

1.4. Brief comments and outline. To the authors' best knowledge, Theorem 1.1 is the only mathematical result claiming Hölder stable determination of unknown coefficients depending on all variables of a partial differential equation (PDE). In the framework of evolution PDEs, where the corresponding inverse problem is to stably retrieve space- and time-varying coefficients, only logarithmic type stability inequalities are available in the mathematical literature, see e.g. [11, 12, 17, 18]. Moreover, it was proved in [39] that logarithmic stable recovery of unknown coefficients of an elliptic PDE is the best one can expect from the DN map, and we refer to [1, 44] for the proof of logarithmic stable determination of the electromagnetic coefficients of the stationary Schrödinger operator.

We point out that upgrading the reconstruction of unknown coefficients from logarithmic-stable to Hölder-stable is interesting not only from a mathematical perspective, but from a numerical and computational perspective as well. Actually, this is all the more important because logarithmic stability can be damaging for the numerical solution of many inverse problems, see e.g. [25, Section 10]. Moreover, even if the identification of unknown coefficients depending on both time- and space-variable is of great interest in its own, it is worth mentioning that it can also be linked with the inverse problem of determining a nonlinear perturbation of a PDE. As a matter of fact it was proved in [17, 29] by mean of a linearization process that the semilinear term entering a nonlinear parabolic equation can be identified by solving the inverse problem of determining the time-dependent coefficient of a related linear parabolic equation. From this viewpoint there is no doubt that Theorem 1.1 is a useful tool for adapting this strategy to the case of semilinear Schrödinger equations.

The remaining part of this article is organized as follows. In the coming section, Section 2, we study the well-posedness of problem (1.1) and we prove that the DN map (1.3) is well defined as a linear bounded operator from $\mathcal{H}(\Sigma)$ into $L^2(\Sigma)$. In Section 3 we build a set of geometrical optics solutions to (1.1) which are the main tool for

deriving Theorem 1.1. Finally, the proof of the stability estimate (1.7) is presented in Section 4, whereas the one of (1.8) is given in Section 5.

2. ANALYSIS OF THE FORWARD PROBLEM

The main result of this section is the following existence and uniqueness result for the IBVP (1.1).

Proposition 2.1. *For $M \in (0, +\infty)$, let $A \in W^{2,\infty}(Q; \mathbb{R})^n$ and $q \in W^{1,\infty}(Q; \mathbb{R})$ satisfy*

$$\|A\|_{W^{2,\infty}(Q)^n} + \|q\|_{W^{1,\infty}(Q)} \leq M. \quad (2.1)$$

Then, for every $g \in \mathcal{H}(\Sigma)$, the system (1.1) admits a unique solution $u \in H^{1,2}(Q)$. Moreover, there exists a positive constant C , depending only on M , T and Ω , such that we have

$$\|u\|_{H^{1,2}(Q)} \leq C \|g\|_{\mathcal{H}(\Sigma)}. \quad (2.2)$$

With reference to (1.3) and the continuity of the trace operator from $H^{1,2}(Q)$ into $L^2(\Sigma)$, Proposition 2.1 immediately entails the following:

Corollary 2.1. *Under the conditions of Proposition 2.1, the DN map $\Lambda_{A,q}$ is well defined by (1.3) and acts as a bounded operator from $\mathcal{H}(\Sigma)$ into $L^2(\Sigma)$.*

The proof of Proposition 2.1 can be found in Section 2.3 by mean of an existence and uniqueness result for the IBVP (1.1) with homogeneous Dirichlet boundary condition and suitable source term, stated in Section 2.2. As a preamble, we recall that the sesquilinear form associated with the operator $-\Delta_{A(t,\cdot)} + q(t,\cdot) = -(\nabla + iA(t,\cdot)) \cdot (\nabla + iA(t,\cdot)) + q(t,\cdot)$ is $H^1(\Omega)$ -elliptic with respect to $L^2(\Omega)$, uniform in $t \in (0, T)$.

2.1. $H^1(\Omega)$ -ellipticity with respect to $L^2(\Omega)$. We define the magnetic gradient ∇_A , associated with $A \in L^\infty(Q)^n$, by

$$\nabla_A u(t, x) := (\nabla + iA(t, x))u(x), \quad u \in H_0^1(\Omega), \quad (t, x) \in Q, \quad (2.3)$$

and for $q \in L^\infty(Q)$, we introduce the sesquilinear form

$$a(t; u, v) := \int_{\Omega} \nabla_A u(t, x) \cdot \overline{\nabla_A v(t, x)} dx + \int_{\Omega} q(t, x) u(x) \overline{v(x)} dx, \quad u, v \in H_0^1(\Omega). \quad (2.4)$$

Then, the Hölder inequality yields

$$\|\nabla_A u(t, \cdot)\|_{L^2(\Omega)^n}^2 \geq \frac{\|\nabla u\|_{L^2(\Omega)^n}^2}{2} - 2\|A\|_{L^\infty(Q)^n}^2 \|u\|_{L^2(\Omega)}^2,$$

for every $u \in H_0^1(\Omega)$ and $t \in (0, T)$, so we get

$$a(t; u, u) + \lambda \|u\|_{L^2(\Omega)}^2 \geq \alpha \|u\|_{H^1(\Omega)}^2, \quad u \in H_0^1(\Omega), \quad t \in (0, T), \quad (2.5)$$

with $\lambda := \frac{1}{2} + \|q\|_{L^\infty(Q)}^2 + 2\|A\|_{L^\infty(Q)^n}^2$ and $\alpha := \frac{1}{2}$.

2.2. Existence and uniqueness result. The proof of Proposition 2.1 essentially boils down to the following existence and uniqueness result for the following IBVP associated with a suitable source term F :

$$\begin{cases} (i\partial_t + \Delta_A + q)v = F & \text{in } Q, \\ v(0, \cdot) = 0 & \text{in } \Omega, \\ v = 0 & \text{on } \Sigma. \end{cases} \quad (2.6)$$

Lemma 2.2. *Let M , A and q be the same as in Proposition 2.1 and let $F \in H^1(0, T; L^2(\Omega))$ verify $F(0, \cdot) = 0$ a.e. in Ω . Then, the system (2.6) admits a unique solution $v \in \mathcal{C}([0, T], H_0^1(\Omega) \cap H^2(\Omega)) \cap \mathcal{C}^1([0, T], L^2(\Omega))$ satisfying*

$$\|v\|_{\mathcal{C}([0, T], H^2(\Omega))} + \|v\|_{\mathcal{C}^1([0, T], L^2(\Omega))} \leq C \|F\|_{H^1(0, T; L^2(\Omega))}, \quad (2.7)$$

for some positive constant C depending only on T , Ω and M .

Proof. We proceed as in the derivation of [37, Section 3, Theorem 10.1] by applying the Faedo-Galerkin method. Namely, we pick a Hilbert basis $\{e_k, k \in \mathbb{N}^*\}$ of $H_0^1(\Omega)$ and consider an approximated solution of size $m \in \mathbb{N}^* := \{1, 2, \dots\}$ of (2.6), of the form

$$v_m(t, x) := \sum_{k=1}^m g_{k,m}(t) e_k(x), \quad (t, x) \in Q, \quad (2.8)$$

where the functions $g_{k,m}$ are defined in such a way that we have

$$\begin{cases} i \langle \partial_t v_m(t, \cdot), e_k \rangle_{L^2(\Omega)} - a(t; v_m(t, \cdot), e_k) = \langle F(t, \cdot), e_k \rangle_{L^2(\Omega)}, & t \in (0, T), \\ v_m(0, \cdot) = 0, \end{cases} \quad (2.9)$$

for all $k = 1, \dots, m$. Since $F \in W^{1,1}(0, T; L^2(\Omega))$ then (2.9) admits a unique solution $v_m \in W^{1,\infty}(0, T; H_0^1(\Omega))$ such that the function $w_m := \partial_t v_m$ solves the following Cauchy problem for every $k = 1, \dots, m$:

$$\begin{cases} i \langle \partial_t w_m(t, \cdot), e_k \rangle_{L^2(\Omega)} - a(t; w_m(t, \cdot), e_k) = a'(t; v_m(t, \cdot), e_k) + \langle \partial_t F(t, \cdot), e_k \rangle_{L^2(\Omega)}, & t \in (0, T), \\ w_m(0) = 0. \end{cases} \quad (2.10)$$

Here, for all $t \in (0, T)$ and all $u, v \in H_0^1(\Omega)$, we have set with reference to (2.3)-(2.4)

$$a'(t; u, v) := i \int_{\Omega} \partial_t A(t, x) \cdot \left(u(x) \overline{\nabla_A v(t, x)} - \nabla_A u(t, x) \overline{v(x)} \right) dx + \int_{\Omega} \partial_t q(t, x) u(x) \overline{v(x)} dx. \quad (2.11)$$

1) The first part of the proof is to establish three *a priori* estimates for the functions v_m and w_m .

a) To this end, we fix $t \in (0, T)$ and we multiply for each $k \in \{1, \dots, m\}$ the first line of (2.9) by $\overline{g_{k,m}(t)}$, sum up over $k = 1, \dots, m$, and infer from (2.8) that

$$i \langle \partial_t v_m(t, \cdot), v_m(t, \cdot) \rangle_{L^2(\Omega)} - a(t; v_m(t, \cdot), v_m(t, \cdot)) = \langle F(t, \cdot), v_m(t, \cdot) \rangle_{L^2(\Omega)}.$$

Taking the imaginary part of both sides of the above identity then yields

$$\frac{d}{ds} \|v_m(s, \cdot)\|_{L^2(\Omega)}^2 = 2 \operatorname{Im} \langle F(s, \cdot), \partial_t v_m(s, \cdot) \rangle_{L^2(\Omega)}, \quad s \in (0, T).$$

Since $v_m(0, \cdot) = 0$, we get upon integrating the above identity over $(0, t)$ that

$$\|v_m(t, \cdot)\|_{L^2(\Omega)}^2 = 2 \operatorname{Im} \int_0^t \langle F(s, \cdot), v_m(s, \cdot) \rangle_{L^2(\Omega)} ds \leq \int_0^t \|F(s, \cdot)\|_{L^2(\Omega)}^2 ds + \int_0^t \|v_m(s, \cdot)\|_{L^2(\Omega)}^2 ds.$$

Therefore, by Gronwall's lemma, there exists a positive constant c_0 , depending only on T, Ω and M such that we have:

$$\|v_m\|_{L^\infty(0, T; L^2(\Omega))} \leq c_0 \|F\|_{L^2(Q)}. \quad (2.12)$$

b) Similarly, by multiplying the first line of (2.9) by $\overline{g'_{k,m}(t)}$, summing up over $k = 1, \dots, m$, and applying (2.8) once more, we get that

$$i \|\partial_t v_m(t, \cdot)\|_{L^2(\Omega)}^2 - a(t; v_m(t, \cdot), \partial_t v_m(t, \cdot)) = \langle F(t, \cdot), \partial_t v_m(t, \cdot) \rangle_{L^2(\Omega)}, \quad t \in (0, T).$$

Upon taking this time the real part in the above identity, we find that

$$a(s; v_m(s, \cdot), \partial_t v_m(s, \cdot)) + a(s; \partial_t v_m(s, \cdot), v_m(s, \cdot)) = -2 \operatorname{Re} \langle F(s, \cdot), \partial_t v_m(s, \cdot) \rangle_{L^2(\Omega)}, \quad s \in (0, T),$$

which may be equivalently rewritten as

$$\frac{d}{ds} a(s; v_m(s, \cdot), v_m(s, \cdot)) = a'(s; v_m(s, \cdot), v_m(s, \cdot)) - 2 \operatorname{Re} \langle F(s, \cdot), \partial_t v_m(s, \cdot) \rangle_{L^2(\Omega)}, \quad s \in (0, T).$$

Now, by integrating with respect to s over $(0, t)$, we obtain that

$$a(t; v_m(t, \cdot), v_m(t, \cdot)) = \int_0^t a'(s; v_m(s, \cdot), v_m(s, \cdot)) ds - 2 \operatorname{Re} \int_0^t \langle F(s, \cdot), \partial_t v_m(s, \cdot) \rangle_{L^2(\Omega)} ds. \quad (2.13)$$

Next, as $\int_0^t \langle F(s, \cdot), \partial_t v_m(s, \cdot) \rangle_{L^2(\Omega)} ds = \langle F(t, \cdot), v_m(t, \cdot) \rangle_{L^2(\Omega)} - \int_0^t \langle \partial_t F(s, \cdot), v_m(s, \cdot) \rangle_{L^2(\Omega)} ds$, we get

$$\left| \int_0^t \langle F(s, \cdot), \partial_t v_m(s, \cdot) \rangle_{L^2(\Omega)} ds \right| \leq \|F(t, \cdot)\|_{L^2(\Omega)} \|v_m(t, \cdot)\|_{L^2(\Omega)} + \int_0^t \|\partial_t F(s, \cdot)\|_{L^2(\Omega)} \|v_m(s, \cdot)\|_{L^2(\Omega)} ds,$$

so that $\|v_m(t, \cdot)\|_{H^1(\Omega)}^2$ can be upper bounded with the aid of (2.5), (2.11) and (2.13), by the following expression

$$c \left(\int_0^t \|v_m(s, \cdot)\|_{H^1(\Omega)}^2 ds + \int_0^t \|\partial_t F(s, \cdot)\|_{L^2(\Omega)}^2 ds \right) + \frac{2}{\alpha} \|F(t, \cdot)\|_{L^2(\Omega)} \|v_m(t, \cdot)\|_{L^2(\Omega)} + \frac{\lambda}{\alpha} \|v_m(t, \cdot)\|_{L^2(\Omega)}^2,$$

where c is a positive constant depending only on M . From this, the two basic inequalities

$$\|F(t, \cdot)\|_{L^2(\Omega)} \|v_m(t, \cdot)\|_{L^2(\Omega)} \leq \frac{\alpha}{4} \|v_m(t, \cdot)\|_{L^2(\Omega)}^2 + \frac{1}{\alpha} \|F(t, \cdot)\|_{L^2(\Omega)}^2$$

and

$$\|F(t, \cdot)\|_{L^2(\Omega)}^2 = 2\text{Re} \int_0^t \langle F(s, \cdot), \partial_t F(s, \cdot) \rangle_{L^2(\Omega)} ds \leq \int_0^t \left(\|F(s, \cdot)\|_{L^2(\Omega)}^2 + \|\partial_t F(s, \cdot)\|_{L^2(\Omega)}^2 \right) ds,$$

and from the estimate (2.12), it then follows that

$$\|v_m(t, \cdot)\|_{H^1(\Omega)}^2 \leq 2c \int_0^t \|v_m(s, \cdot)\|_{H^1(\Omega)}^2 ds + 2 \left(c + \frac{2}{\alpha^2} + \frac{c_0^2}{4} + \frac{\lambda c_0^2}{\alpha} \right) \|F\|_{H^1(0, T; L^2(\Omega))}^2.$$

Then, an application of Gronwall's lemma provides a constant $C = C(T, M, \alpha) \in (0, +\infty)$ such that

$$\|v_m\|_{L^\infty(0, T; H^1(\Omega))} \leq C \|F\|_{H^1(0, T; L^2(\Omega))}. \quad (2.14)$$

c) Further, we put $p(t, x) := |A(t, x)|^2 + q(t, x)$ for $(t, x) \in Q$, integrate by parts in the first integral of (2.11) and obtain for all $u, v \in H_0^1(\Omega)$ that

$$\begin{aligned} a'(t; u, v) &= i \int_{\Omega} \left(u(x) \partial_t A(t, x) \cdot \overline{\nabla v(x)} - \partial_t A(t, x) \cdot \nabla u(x) \overline{v(x)} \right) dx + \int_{\Omega} \partial_t p(t, x) u(x) \overline{v(x)} dx \\ &= \int_{\Omega} \left(-2i \partial_t A(t, x) \cdot \nabla u(x) + (\partial_t p(t, x) - i \nabla \cdot \partial_t A(t, x)) u(x) \right) \overline{v(x)} dx. \end{aligned}$$

This and (2.10) yield

$$\begin{cases} i \langle \partial_t w_m(t, \cdot), e_k \rangle_{L^2(\Omega)} - a(t; w_m(t, \cdot), e_k) = \langle F_m(t, \cdot), e_k \rangle_{L^2(\Omega)}, & t \in (0, T), \\ w_m(0) = 0, \end{cases} \quad (2.15)$$

for all $k = 1, \dots, m$, where

$$F_m(t, x) := -2i \partial_t A(t, x) \cdot \nabla v_m(t, x) + (\partial_t p(t, x) - i \nabla \cdot \partial_t A(t, x)) v_m(t, x) + \partial_t F(t, x), \quad (t, x) \in Q. \quad (2.16)$$

Next, multiplying the first line in (2.15) by $g'_{k,m}(t)$ and summing up over $k = 1, \dots, m$, lead to

$$i \langle \partial_t w_m(t, \cdot), w_m(t, \cdot) \rangle_{L^2(\Omega)} - a(t; w_m(t, \cdot), w_m(t, \cdot)) = \langle F_m(t, \cdot), w_m(t, \cdot) \rangle_{L^2(\Omega)}, \quad t \in (0, T).$$

Therefore, we have

$$\frac{d}{ds} \|w_m(s, \cdot)\|_{L^2(\Omega)}^2 = 2\text{Im} \langle F_m(s, \cdot), w_m(s, \cdot) \rangle_{L^2(\Omega)}, \quad s \in (0, T).$$

Now, for each $t \in (0, T)$, we find upon integrating both sides of the above identity over $(0, t)$ that

$$\|w_m(t, \cdot)\|_{L^2(\Omega)}^2 \leq \int_0^t \|w_m(s, \cdot)\|_{L^2(\Omega)}^2 ds + \int_0^t \|F_m(s, \cdot)\|_{L^2(\Omega)}^2 ds,$$

which, combined with (2.16), entails

$$\|w_m(t, \cdot)\|_{L^2(\Omega)}^2 \leq \int_0^t \|w_m(s, \cdot)\|_{L^2(\Omega)}^2 ds + c \left(\|\partial_t F\|_{L^2(Q)}^2 + \|v_m\|_{L^\infty(0, T; H^1(\Omega))}^2 \right),$$

for some constant $c = c(T, M) \in (0, +\infty)$. Therefore, we have

$$\|w_m(t, \cdot)\|_{L^2(\Omega)}^2 \leq c \left(\|\partial_t F\|_{L^2(Q)}^2 + \|v_m\|_{L^\infty(0, T; H^1(\Omega))}^2 \right), \quad t \in (0, T),$$

by Gronwall's lemma, and consequently

$$\|w_m\|_{L^\infty(0,T;L^2(\Omega))} \leq C\|F\|_{H^1(0,T;L^2(\Omega))}, \quad (2.17)$$

by (2.14), where C is another positive constant depending only on T , M and α .

2) Having established (2.14) and (2.17), we turn now to showing existence of a solution to (2.6). This can be done in accordance with (2.14) by extracting a subsequence $(v_{m'})_{m'}$ of $(v_m)_m$, which converges to $v \in L^\infty(0, T; H_0^1(\Omega))$ in the weak-star topology of $L^\infty(0, T; H_0^1(\Omega))$. By substituting m' for m in (2.9) and sending m' to infinity, we get that

$$\begin{cases} (i\partial_t + \Delta_A + q)v = F & \text{in } Q, \\ v(0, \cdot) = 0 & \text{in } \Omega. \end{cases} \quad (2.18)$$

As a consequence, we have $\partial_t v \in L^\infty(0, T; H^{-1}(\Omega))$ and hence $v \in L^\infty(0, T; H_0^1(\Omega)) \cap W^{1,\infty}(0, T; H^{-1}(\Omega))$. Further, due to (2.17) and the Banach-Alaoglu theorem, there exists a subsequence of $(w_m)_m$ which converges to $w \in L^\infty(0, T; L^2(\Omega))$ in the weak-star topology of $L^\infty(0, T; L^2(\Omega))$. Since $w_m = \partial_t v_m$ for every $m \in \mathbb{N}^*$ then we necessarily have $\partial_t v = w \in L^\infty(0, T; L^2(\Omega))$ and thus $v \in L^\infty(0, T; H_0^1(\Omega)) \cap W^{1,\infty}(0, T; L^2(\Omega))$. Further, by arguing in the exact same way as in the derivation of [37, Theorem 8.3 and Remark 10.2, Chapter 3], we get that $v \in \mathcal{C}([0, T]; H_0^1(\Omega)) \cap \mathcal{C}^1([0, T]; L^2(\Omega))$. Moreover, for all fixed $t \in [0, T]$, we deduce from (2.18) that $v(t, \cdot)$ is solution to the elliptic boundary value problem

$$\begin{cases} (\Delta_A + q(t, \cdot))v(t, \cdot) = F(t, \cdot) - i\partial_t v(t, \cdot) & \text{in } \Omega, \\ v(t, \cdot) = 0, & \text{on } \partial\Omega. \end{cases}$$

As $F - i\partial_t v \in \mathcal{C}([0, T]; L^2(\Omega))$ then we have $v \in \mathcal{C}([0, T], H_0^1(\Omega) \cap H^2(\Omega))$ and (2.7) follows directly from this, (2.4), (2.14) and (2.17). \square

Remark 2.3. a) With reference to (2.12), we point out for further use that the solution v to (2.2) satisfies the estimate

$$\|v\|_{L^\infty(0,T;L^2(\Omega))} \leq c_0\|F\|_{L^2(Q)},$$

for some constant c_0 depending only on Ω , T and M .

b) Let M , A and q be the same as in Lemma 2.2, and let $F \in H^1(0, T; L^2(\Omega))$ satisfy $F(T, \cdot) = 0$. Putting $\tilde{A}(t, x) := -A(T - t, x)$, $\tilde{q}(t, x) := q(T - t, x)$ and $\tilde{F}(t, x) := F(T - t, x)$ for $(t, x) \in Q$, we see that we have

$$\begin{cases} (i\partial_t + \Delta_A + q)v = F & \text{in } Q, \\ v(T, \cdot) = 0 & \text{in } \Omega, \\ v = 0 & \text{on } \Sigma, \end{cases} \quad (2.19)$$

if and only if $\tilde{v}(t, x) := \overline{v(T - t, x)}$ is a solution to the system (2.6) where $(\tilde{A}, \tilde{q}, \tilde{F})$ is substituted for (A, q, F) . Therefore, by Lemma 2.2, there exists a unique solution $v \in \mathcal{C}([0, T], H_0^1(\Omega) \cap H^2(\Omega)) \cap \mathcal{C}^1([0, T], L^2(\Omega))$ to (2.19), and it is clear that v verifies the estimate (2.7).

2.3. Completion of the proof of Proposition 2.1. In light of [38, Theorem 2.3, Chapter 4] there exists $G \in H^3(Q)$ satisfying

$$G(0, \cdot) = \partial_t G(0, \cdot) = 0 \text{ in } \Omega \text{ and } G = g \text{ on } \Gamma,$$

and

$$\|G\|_{H^3(Q)} \leq C\|g\|_{\mathcal{H}(\Sigma)}, \quad (2.20)$$

for some positive constant C , depending only on Ω and T . Therefore, the function

$$F := -(i\partial_t + \Delta_A + q)G \in H^1(0, T; L^2(\Omega)) \quad (2.21)$$

verifies $F(0, \cdot) = 0$ in Ω . Let v be the $\mathcal{C}([0, T], H_0^1(\Omega) \cap H^2(\Omega)) \cap \mathcal{C}^1([0, T], L^2(\Omega))$ -solution to (2.6), associated with the source term F defined by (2.21), which is given by Lemma 2.2. Then, $u := G + v \in H^{1,2}(Q)$ is a solution to

(1.1) and (2.2) follows directly from (2.7) and (2.20). Finally, we get that such a solution is unique by applying (2.2) with $g = 0$.

3. GO SOLUTIONS

In this section we build appropriate geometric optics (GO) solutions to the magnetic Schrödinger equation in Q , which are used in the derivation of the stability estimates of Theorem 1.1, presented in Sections 4 and 5.

Namely, given $A_j \in W^{5,\infty}(Q)^n \cap H^6(Q)^n$ and $q_j \in W^{4,\infty}(Q)$, $j = 1, 2$, fulfilling the conditions (1.4)-(1.6), we seek a solution u_j to the magnetic Schrödinger equation

$$(i\partial_t + \Delta_{A_j} + q_j)u_j = 0 \text{ in } Q, \quad (3.1)$$

of the form

$$u_j(t, x) = \varphi_\sigma(t, x) (u_{j,1}(t, x) + \sigma^{-1}u_{j,2}(t, x)) + r_{j,\sigma}(t, x) \text{ with } \varphi_\sigma(t, x) := e^{i\sigma(-\sigma t + x \cdot \omega)}. \quad (3.2)$$

Here $\sigma \in (1, +\infty)$ and $\omega \in \mathbb{S}^{n-1} := \{y \in \mathbb{R}^n; |y| = 1\}$ are arbitrarily fixed and the remainder term $r_{j,\sigma}$ in the asymptotic expansion of u_j with respect to σ^{-1} , scales at most like σ^{-1} as $\sigma \rightarrow +\infty$, in a sense that we will make precise below. Moreover, we impose that $u_{j,1}$ and $u_{j,2}$ be in $H^3(Q)$, and that they satisfy

$$\omega \cdot \nabla_{A_j} u_{j,1} = 0 \text{ and } 2i\omega \cdot \nabla_{A_j} u_{j,2} + (i\partial_t + \Delta_{A_j} + q_j)u_{j,1} = 0 \text{ in } Q, \quad (3.3)$$

and

$$\exists \delta \in \left(0, \frac{T}{4}\right), (t \in (0, \delta) \cup (T - \delta, T)) \Rightarrow (u_{j,1}(t, x) = u_{j,2}(t, x) = 0, x \in \Omega). \quad (3.4)$$

The two conditons in (3.3) can be understood from the formal commutator formula $[i\partial_t + \Delta_{A_j}, \varphi_\sigma] = i\partial_t \varphi_\sigma + \Delta \varphi_\sigma + 2\nabla \varphi_\sigma \cdot \nabla_{A_j} = 2i\sigma \varphi_\sigma \omega \cdot \nabla_{A_j}$, entailing

$$\begin{aligned} (i\partial_t + \Delta_{A_j} + q_j)(u_j - r_{j,\sigma}) &= (i\partial_t + \Delta_{A_j} + q_j)\varphi_\sigma(u_{j,1} + \sigma^{-1}u_{j,2}) \\ &= \varphi_\sigma(2i\sigma\omega \cdot \nabla_{A_j} u_{j,1} + (i\partial_t + \Delta_{A_j} + q_j)u_{j,1} + 2i\omega \cdot \nabla_{A_j} u_{j,2} + \sigma^{-1}(i\partial_t + \Delta_{A_j} + q_j)u_{j,2}) \end{aligned}$$

in Q . This and (3.1)-(3.2) lead to defining $r_{1,\sigma}$ by

$$\begin{cases} (i\partial_t + \Delta_{A_1} + q_1)r_{1,\sigma} = -\sigma^{-1}\varphi_\sigma(i\partial_t + \Delta_{A_1} + q_1)u_{1,2} & \text{in } Q, \\ r_{1,\sigma}(0, \cdot) = 0 & \text{in } \Omega, \\ r_{1,\sigma} = 0 & \text{on } \Sigma, \end{cases} \quad (3.5)$$

and $r_{2,\sigma}$ by

$$\begin{cases} (i\partial_t + \Delta_{A_2} + q_2)r_{2,\sigma} = -\sigma^{-1}\varphi_\sigma(i\partial_t + \Delta_{A_2} + q_2)u_{2,2} & \text{in } Q, \\ r_{2,\sigma}(T, \cdot) = 0 & \text{in } \Omega, \\ r_{2,\sigma} = 0 & \text{on } \Sigma. \end{cases} \quad (3.6)$$

The initial condition in (3.5) and the final condition in (3.6) are imposed in such a way that the product $r_{1,\sigma}\overline{r_{2,\sigma}}$ vanishes at both ends of the time interval $(0, T)$.

The first step of the construction of the functions $u_{j,k}$, for $j, k = 1, 2$, involves extending the two magnetic potentials A_1 and A_2 to $(0, T) \times \mathbb{R}^n$ as follows. First, we refer to [43, Theorem 5 in Section 3] and pick a magnetic potential $\tilde{A}_1 \in W^{5,\infty}((0, T) \times \mathbb{R}^n; \mathbb{R}^n) \cap H^6((0, T) \times \mathbb{R}^n; \mathbb{R}^n)$ which coincides with A_1 in Q and satisfies

$$\exists R \in (0, +\infty), \forall t \in [0, T], \text{supp } \tilde{A}_1(t, \cdot) \subset \{x \in \mathbb{R}^n, |x| \leq R\}$$

and the estimate

$$\|\tilde{A}_1\|_{W^{5,\infty}((0,T) \times \mathbb{R}^n)^n} \leq C\|A_1\|_{W^{5,\infty}(Q)^n} \text{ and } \|\tilde{A}_1\|_{H^6((0,T) \times \mathbb{R}^n)^n} \leq C\|A_1\|_{H^6(Q)^n}, \quad (3.7)$$

for some positive constant C depending only on T and Ω . Thus, putting

$$\tilde{A}_2(t, x) := \begin{cases} A_2(t, x) & \text{if } x \in \Omega, \\ \tilde{A}_1(t, x) & \text{if } x \in \mathbb{R}^n \setminus \Omega, \end{cases} \quad (3.8)$$

we infer from (1.4) that $\tilde{A}_2 \in W^{5,\infty}((0, T) \times \mathbb{R}^n)^n \cap H^6((0, T) \times \mathbb{R}^n)^n$. Moreover, it is clear from (3.7)-(3.8) upon possibly substituting $\max(1, C)$ for C in (3.7), that

$$\|\tilde{A}_j\|_{W^{5,\infty}((0,T)\times\mathbb{R}^n)^n} \leq C \max_{k=1,2} \|A_k\|_{W^{5,\infty}(Q)^n} \text{ and } \|\tilde{A}_j\|_{H^6((0,T)\times\mathbb{R}^n)^n} \leq C \max_{k=1,2} \|A_k\|_{H^6(Q)^n}, \quad j = 1, 2. \quad (3.9)$$

The next step is to introduce two functions, the first one $\chi = \chi_\delta \in C^\infty(\mathbb{R}; [0, 1])$, being supported in $(\delta, T - \delta)$, satisfies $\chi(t) = 1$ if $t \in [2\delta, T - 2\delta]$ and fulfills

$$\forall k \in \mathbb{N}, \exists C_k \in (0, +\infty), \|\chi\|_{W^{k,\infty}(\mathbb{R})} \leq C_k \delta^{-k},$$

whereas the second one is defined for $\tau \in \mathbb{R}, \xi \in \omega^\perp := \{x \in \mathbb{R}^n : x \cdot \omega = 0\}$ and $y \in \mathbb{S}^{n-1} \cap \omega^\perp$, by

$$\beta(t, x) := y \cdot \nabla \left(e^{-i(t\tau + \xi \cdot x)} \exp \left(-i \int_{\mathbb{R}} A(t, x + s\omega) \cdot \omega ds \right) \right), \quad (t, x) \in (0, T) \times \mathbb{R}^n. \quad (3.10)$$

Here we have set $A := \tilde{A}_1 - \tilde{A}_2$ (that is $A = A_1 - A_2$ in Q and $A = 0$ in $((0, T) \times \mathbb{R}^n) \setminus Q$) in such a way that we have $\omega \cdot \nabla \beta(t, x) = 0$ for all $(t, x) \in (0, T) \times \mathbb{R}^n$.

Now, a direct calculation shows that each of the two functions

$$u_{1,1}(t, x) := \chi(t)\beta(t, x) \exp \left(i \int_0^{+\infty} \tilde{A}_1(t, x + s\omega) \cdot \omega ds \right)$$

and

$$u_{2,1}(t, x) := \chi(t) \exp \left(i \int_0^{+\infty} \tilde{A}_2(t, x + s\omega) \cdot \omega ds \right)$$

is a solution to the first equation of (3.3) satisfying the condition (3.4). Further, it follows from this, (1.6) and (3.9)-(3.10) that

$$\|u_{1,1}\|_{H^3(Q)} \leq C \langle \tau, \xi \rangle^4 \delta^{-3} \text{ and } \|u_{2,1}\|_{H^3(Q)} \leq C \delta^{-3}, \quad (3.11)$$

where C denotes a positive constant depending only Ω, T and M , which may change from line to line, and $\langle \tau, \xi \rangle$ is a shorthand for $(1 + \tau^2 + \xi^2)^{1/2}$.

Similarly, using that any $x \in \mathbb{R}^n$ decomposes into the sum $x = x_\perp + s\omega$ with $s := x \cdot \omega$ and $x_\perp := x - s\omega \in \omega^\perp$, it can be checked through standard computations that

$$u_{j,2}(t, x_\perp + s\omega) := -\frac{1}{2i} \int_0^s \exp \left(-i \int_{s_1}^s \tilde{A}_j(t, x_\perp + s_2\omega) \cdot \omega ds_2 \right) (i\partial_t + \Delta_{A_j} + q_j) u_{j,1}(t, x_\perp + s_1\omega) ds_1, \quad j = 1, 2,$$

is a solution to the second equation of (3.3) obeying the condition (3.4). Further, by (1.6) and (3.9) we have

$$\|u_{1,2}\|_{H^3(Q)} \leq C \langle \tau, \xi \rangle^6 \delta^{-4} \text{ and } \|u_{2,2}\|_{H^3(Q)} \leq C \delta^{-4}, \quad (3.12)$$

and

$$\|(i\partial_t + \Delta_{A_1} + q_1)u_{1,2}\|_{L^2(Q)} \leq C \langle \tau, \xi \rangle^5 \delta^{-2} \text{ and } \|(i\partial_t + \Delta_{A_2} + q_2)u_{2,2}\|_{L^2(Q)} \leq C \delta^{-2}. \quad (3.13)$$

Having specified $u_{j,k}$ for $j, k = 1, 2$, we turn now to examining the remainder term $r_{j,\sigma}$. We first infer from Lemma 2.2 (resp., Statement b) of Remark 2.3) that $r_{1,\sigma}$ (resp. $r_{2,\sigma}$) is well defined as the $\mathcal{C}([0, T], H_0^1(\Omega) \cap H^2(\Omega)) \cap C^1([0, T], L^2(\Omega))$ -solution to (3.5) (resp., (3.6)). Next, Statement a) in Remark 2.3 and (3.13) yield

$$\|r_{1,\sigma}\|_{L^2(Q)} \leq C \|(i\partial_t + \Delta_{A_1} + q_1)u_{1,2}\|_{L^2(Q)} \sigma^{-1} \leq C \langle \tau, \xi \rangle^5 \delta^{-2} \sigma^{-1}. \quad (3.14)$$

On the other hand, we know from (2.7) and (3.12) that

$$\|r_{1,\sigma}\|_{L^2(0,T;H^2(\Omega))} \leq C \|e^{i\sigma(-\sigma t + x \cdot \omega)} (i\partial_t + \Delta_{A_1} + q_1)u_{1,2}\|_{H^1(0,T;L^2(\Omega))} \sigma^{-1} \leq C \langle \tau, \xi \rangle^6 \delta^{-3} \sigma,$$

Thus, interpolating with (3.14), we have $\|r_{1,\sigma}\|_{L^2(0,T;H^1(\Omega))} \leq C \langle \tau, \xi \rangle^6 \delta^{-3}$ and hence

$$\|r_{1,\sigma}\|_{L^2(0,T;H^1(\Omega))} + \sigma \|r_{1,\sigma}\|_{L^2(Q)} \leq C \langle \tau, \xi \rangle^6 \delta^{-3}. \quad (3.15)$$

Analogously, we establish that

$$\|r_{2,\sigma}\|_{L^2(0,T;H^1(\Omega))} + \sigma \|r_{2,\sigma}\|_{L^2(Q)} \leq C \delta^{-3}. \quad (3.16)$$

Having built $u_{j,k}$ and $r_{j,\sigma}$, for $j, k = 1, 2$, fulfilling (3.1)–(3.4), we are now in position to derive the stability estimates (1.7)–(1.8) of Theorem 1.1.

4. PROOF OF THE STABILITY ESTIMATE (1.7)

We stick to the notations of Section 3 and recall from (3.4) that

$$u_1(0, x) = u_2(T, x) = 0, \quad x \in \Omega. \quad (4.1)$$

The proof of (1.7) boils down to a suitable estimate of the Fourier transform of the function $\chi^2 A$, presented in Lemma 4.2.

4.1. Estimation of the Fourier transform of $\chi^2 A$. We start by proving the following technical estimate.

Lemma 4.1. *There exists a constant $C = C(T, \Omega, M) \in (0, +\infty)$ such that we have*

$$\begin{aligned} & \left| \int_{\mathbb{R}^{n+1}} \chi^2(t) \beta(t, x) e^{i \int_0^{+\infty} A(t, x+s\omega) \cdot \omega ds} A(t, x) \cdot \omega dx dt \right| \\ & \leq C (\langle \tau, \xi \rangle^6 \delta^{-8} \sigma^5 \|\Lambda_{A_1, q_1} - \Lambda_{A_2, q_2}\| + \langle \tau, \xi \rangle^8 \delta^{-6} \sigma^{-1}), \end{aligned} \quad (4.2)$$

uniformly in $\xi \in \omega^\perp$.

Proof. For $j = 1, 2$, put $\psi_{j, \sigma} := u_j - r_{j, \sigma} = u_j$ in Σ and let v_2 be the $H^{1,2}(Q)$ -solution to

$$\begin{cases} (i\partial_t + \Delta_{A_2} + q_2)v_2 = 0 & \text{in } Q, \\ v_2(0, \cdot) = 0 & \text{in } \Omega, \\ v_2 = \psi_{1, \sigma} & \text{on } \Sigma, \end{cases}$$

given by Proposition 2.1. In light of (3.1) the function $w := v_2 - u_1$ then solves

$$\begin{cases} (i\partial_t + \Delta_{A_2} + q_2)w = 2iA \cdot \nabla u_1 + V u_1 & \text{in } Q, \\ w(0, \cdot) = 0 & \text{in } \Omega, \\ w = 0 & \text{on } \Sigma, \end{cases}$$

with $V := i\nabla \cdot A - (|A_1|^2 - |A_2|^2) + q_1 - q_2$. Next, by multiplying the first equation of the above system by $\overline{u_2}$ and integrating by parts over Q , we deduce from (3.1) and (4.1) that

$$\int_Q (2iA \cdot \nabla + V)u_1(t, x) \overline{u_2(t, x)} dx dt = \int_\Sigma \partial_\nu w(t, x) \overline{u_2(t, x)} d\sigma(x) dt. \quad (4.3)$$

Further, since $(\partial_\nu + iA_2 \cdot \nu)v_2 = \Lambda_{A_2, q_2} \psi_{1, \sigma}$ and $(\partial_\nu + iA_1 \cdot \nu)u_1 = \Lambda_{A_1, q_1} \psi_{1, \sigma}$, we have $\partial_\nu w = (\Lambda_{A_2, q_2} - \Lambda_{A_1, q_1})\psi_{1, \sigma}$ in virtue of (1.4), and hence

$$\begin{aligned} \left| \int_\Sigma \partial_\nu w(t, x) \overline{u_2(t, x)} d\sigma(x) dt \right| & \leq \|(\Lambda_{A_2, q_2} - \Lambda_{A_1, q_1})\psi_{1, \sigma}\|_{L^2(\Sigma)} \|\psi_{2, \sigma}\|_{L^2(\Sigma)} \\ & \leq C \|\Lambda_{A_1, q_1} - \Lambda_{A_2, q_2}\| \|\psi_{1, \sigma}\|_{\mathcal{H}(\Sigma)} \|\psi_{2, \sigma}\|_{L^2(\Sigma)} \\ & \leq C \|\Lambda_{A_2, q_2} - \Lambda_{A_2, q_2}\| \langle \tau, \xi \rangle^6 \delta^{-8} \sigma^6, \end{aligned} \quad (4.4)$$

by Corollary 2.1, the continuity of the trace operator from $H^3(Q)$ into $\mathcal{H}(\Sigma)$, (3.2) and the estimates (3.11)-(3.12).

On the other hand, we know from (3.2) that

$$\begin{aligned} \int_Q (2iA \cdot \nabla + V)u_1 \overline{u_2}(t, x) dx dt & = -2\sigma \int_Q (A \cdot \omega) u_{1,1} \overline{u_{2,1}}(t, x) dx dt + r \\ & = -2\sigma \int_Q \chi^2(t) \beta(t, x) e^{i \int_0^{+\infty} A(t, x+s\omega) \cdot \omega ds} A(t, x) \cdot \omega dx dt + r, \end{aligned} \quad (4.5)$$

where

$$r := -2\sigma \int_Q u_{1,1} (\overline{u_{2,2}} \sigma^{-1} + \varphi_\sigma \overline{r_{2, \sigma}}) A(t, x) \cdot \omega dx dt$$

$$-2 \int_Q A \cdot (\varphi_\sigma (\nabla u_{1,1} + \nabla u_{1,2} \sigma^{-1}) + \nabla r_{1,\sigma}) \bar{u}_2(t, x) dx dt + \int_Q V u_1 \bar{u}_2(t, x) dx dt.$$

Since $|r| \leq C \langle \tau, \xi \rangle^8 \delta^{-6}$ by (3.15)-(3.16), it then follows from (4.5) that

$$\begin{aligned} & \sigma^{-1} \left| \int_Q (2iA \cdot \nabla + V) u_1 \bar{u}_2(t, x) dx dt \right| \\ & \geq 2 \left| \int_Q \chi^2(t) \beta(t, x) e^{i \int_0^{+\infty} A(t, x+s\omega) \cdot \omega ds} A(t, x) \cdot \omega dx dt \right| - C \langle \tau, \xi \rangle^8 \delta^{-6} \sigma^{-1}, \end{aligned}$$

which, combined with (4.3)-(4.4), yields (4.2). \square

Having established Lemma 4.1 we may now estimate the Fourier transform of χA . We recall that the Fourier transform of a function $f \in L^1(\mathbb{R}^{1+n})^n$ is defined for all $(\tau, \xi) \in \mathbb{R} \times \mathbb{R}^n$ by

$$\hat{f}(\tau, \xi) := (2\pi)^{-\frac{n+1}{2}} \int_{\mathbb{R}^{1+n}} e^{-i(\tau+x \cdot \xi)} f(t, x) dx dt.$$

Lemma 4.2. *There exists a positive constant C , depending only on T , Ω and M , such that the inequality*

$$|\xi| \left| \widehat{\chi^2 A}(\tau, \xi) \right| \leq C \left(\langle \tau, \xi \rangle^6 \delta^{-8} \sigma^5 \|\Lambda_{A_1, q_1} - \Lambda_{A_2, q_2}\| + \langle \tau, \xi \rangle^8 \delta^{-6} \sigma^{-1} \right), \quad (4.6)$$

holds for any $(\tau, \xi) \in \mathbb{R}^{1+n}$.

Proof. The estimate (4.6) being obviously true for $\xi = 0$, we will solely focus on the case $\xi \neq 0$. We fix $\omega \in \xi^\perp \cap \mathbb{S}^{n-1}$ and we use the decomposition $x = x_\perp + \kappa \omega$, where $\kappa := x \cdot \omega$ and $x_\perp := x - \kappa \omega$, and recall from (3.10) that we have $\beta(t, x) = \beta(t, x_\perp)$, so we obtain

$$\begin{aligned} & \int_{\mathbb{R}^{1+n}} \chi^2(t) \beta(t, x) e^{i \int_0^{+\infty} A(t, x+s\omega) \cdot \omega ds} A(t, x) \cdot \omega dx dt \\ &= \int_{\mathbb{R}} \int_{\mathbb{R}} \int_{\omega^\perp} \chi^2(t) \beta(t, x_\perp) e^{i \int_\kappa^{+\infty} A(t, x_\perp+s\omega) \cdot \omega ds} A(t, x_\perp + \kappa \omega) \cdot \omega dx_\perp d\kappa dt \\ &= i \int_{\mathbb{R}} \int_{\omega^\perp} \chi^2(t) \beta(t, x_\perp) \left(\int_{\mathbb{R}} \partial_\kappa e^{i \int_\kappa^{+\infty} A(t, x_\perp+s\omega) \cdot \omega ds} d\kappa \right) dx_\perp dt \\ &= i \int_{\mathbb{R}} \chi^2(t) \left(\int_{\omega^\perp} \beta(t, x_\perp) \left(1 - e^{i \int_{\mathbb{R}} A(t, x_\perp+s\omega) \cdot \omega ds} \right) dx_\perp \right) dt \\ &= i \int_{\mathbb{R}} \chi^2(t) e^{-it\tau} \left(\int_{\omega^\perp} y \cdot \nabla \left(e^{-i\xi \cdot x_\perp} e^{-i \int_{\mathbb{R}} A(t, x_\perp+s\omega) \cdot \omega ds} \right) \left(1 - e^{i \int_{\mathbb{R}} A(t, x_\perp+s\omega) \cdot \omega ds} \right) dx_\perp \right) dt. \quad (4.7) \end{aligned}$$

Next, as we have

$$\begin{aligned} & \int_{\omega^\perp} y \cdot \nabla \left(e^{-i\xi \cdot x_\perp} e^{-i \int_{\mathbb{R}} A(t, x_\perp+s\omega) \cdot \omega ds} \right) \left(1 - e^{i \int_{\mathbb{R}} A(t, x_\perp+s\omega) \cdot \omega ds} \right) dx_\perp \\ &= \int_{\omega^\perp} \nabla \cdot \left(y e^{-i\xi \cdot x_\perp} e^{-i \int_{\mathbb{R}} A(t, x_\perp+s\omega) \cdot \omega ds} \right) \left(1 - e^{i \int_{\mathbb{R}} A(t, x_\perp+s\omega) \cdot \omega ds} \right) dx_\perp \\ &= -i \int_{\omega^\perp} e^{-i\xi \cdot x_\perp} y \cdot \nabla \left(\int_{\mathbb{R}} A(t, x_\perp + s\omega) \cdot \omega ds \right) dx_\perp, \end{aligned}$$

by integrating by parts, (4.7) and the Fubini theorem entail

$$\begin{aligned} & \int_{\mathbb{R}^{1+n}} \chi^2(t) \beta(t, x) e^{i \int_0^{+\infty} A(t, x+s\omega) \cdot \omega ds} A(t, x) \cdot \omega dx dt \\ &= \int_{\mathbb{R}} \chi^2(t) e^{-it\tau} \left(\int_{\omega^\perp} \left(\int_{\mathbb{R}} y \cdot \nabla (A(t, x_\perp + s\omega) \cdot \omega) ds \right) e^{-i\xi \cdot x_\perp} dx_\perp \right) dt \\ &= \int_{\mathbb{R}} \chi^2(t) e^{-it\tau} \left(\int_{\mathbb{R}^n} e^{-i\xi \cdot x} y \cdot \nabla (A(t, x) \cdot \omega) dx \right) dt \end{aligned}$$

$$= \int_{\mathbb{R}^{n+1}} e^{-i(t\tau+x\cdot\xi)} \chi^2(t) y \cdot \nabla(A(t, x) \cdot \omega) dx dt.$$

Therefore, taking $y = \frac{\xi}{|\xi|}$ and applying Stokes formula to the above integral, we obtain

$$\begin{aligned} & i \int_{\mathbb{R}^{1+n}} \chi^2(t) \beta(t, x) e^{i \int_0^{+\infty} A(t, x+s\omega) \cdot \omega ds} A(t, x) \cdot \omega dx dt \\ &= -|\xi| \int_{\mathbb{R}^{n+1}} e^{-i(t\tau+x\cdot\xi)} \chi^2(t) A(t, x) \cdot \omega dx dt \\ &= -(2\pi)^{\frac{n+1}{2}} |\xi| \widehat{\chi^2 A}(\tau, \xi) \cdot \omega, \end{aligned}$$

which yields

$$|\xi| \left| \widehat{\chi^2 A}(\tau, \xi) \cdot \omega \right| \leq C \langle \tau, \xi \rangle^6 \delta^{-8} (\|\Lambda_{A_2, q_2} - \Lambda_{A_2, q_2}\| \sigma^5 + \langle \tau, \xi \rangle^2 \delta^2 \sigma^{-1}), \quad (4.8)$$

in virtue of (4.2).

Further, since $\nabla \cdot A = 0$ in Q , by (1.5), then we have $\widehat{\chi^2 A}(\tau, \xi) \cdot \xi = 0$ by direct calculation and hence

$$\widehat{\chi^2 A}(\tau, \xi) = \sum_{k=1}^{n-1} \left(\widehat{\chi^2 A}(\tau, \xi) \cdot e_k \right) e_k,$$

for any orthonormal basis $\{e_1, \dots, e_{n-1}\}$ of ξ^\perp . Finally, (4.6) follows directly from this upon applying (4.8) with $\omega = e_k$ for $k = 1, \dots, n-1$. \square

Having established Lemma 4.2, we are now in position to derive the stability estimate (1.7).

4.2. Completion of the proof. We start by estimating $\|\chi^2 A\|_{L^2(0, T; H^5(\Omega))^n}$. Recalling from (1.4) that $\chi^2 A$ is supported in $(\delta, T - \delta) \times \overline{\Omega}$, we see that

$$\|\chi^2 A\|_{L^2(0, T; H^5(\Omega))^n}^2 = \int_{\mathbb{R}^{1+n}} \langle \xi \rangle^{10} |\widehat{\chi^2 A}(\tau, \xi)|^2 d\xi d\tau,$$

where, as usual, $\langle \xi \rangle$ denotes $(1 + |\xi|^2)^{\frac{1}{2}}$. Next, for $R \in (1, +\infty)$ fixed, we put $B_R := \{(\tau, \xi) \in \mathbb{R}^{1+n}; |(\tau, \xi)| < R\}$, use that $\chi^2 A \in H^1(\mathbb{R}; H^5(\mathbb{R}^n))^n \cap L^2(\mathbb{R}; H^6(\mathbb{R}^n))^n$, and obtain

$$\begin{aligned} \int_{\mathbb{R}^{1+n} \setminus B_R} \langle \xi \rangle^{10} |\widehat{\chi^2 A}(\tau, \xi)|^2 d\xi d\tau &\leq R^{-2} \int_{\mathbb{R}^{1+n} \setminus B_R} |(\xi, \tau)|^2 \langle \xi \rangle^{10} |\widehat{\chi^2 A}(\tau, \xi)|^2 d\xi d\tau \\ &\leq R^{-2} \left(\|\chi^2 A\|_{H^1(\mathbb{R}; H^5(\mathbb{R}^n))^n}^2 + \|\chi^2 A\|_{L^2(\mathbb{R}; H^6(\mathbb{R}^n))^n}^2 \right). \end{aligned}$$

Therefore, we have

$$\int_{\mathbb{R}^{1+n} \setminus B_R} \langle \xi \rangle^{10} |\widehat{\chi^2 A}(\tau, \xi)|^2 d\xi d\tau \leq CR^{-2} \delta^{-2} \|A\|_{H^6(Q)^n}^2 \leq CR^{-2} \delta^{-2}, \quad (4.9)$$

where C is a generic positive constant depending only on Ω , T and M , which may change from line to line. Further, setting $E_R := \{(\tau, \xi) \in \mathbb{R}^{1+n}; |\xi| \leq R^{-\frac{3}{n}}\}$, we get

$$\begin{aligned} \int_{B_R \cap E_R} \langle \xi \rangle^{10} |\widehat{\chi^2 A}(\tau, \xi)|^2 d\xi d\tau &\leq \int_{-R}^R \int_{|\xi| \leq R^{-\frac{3}{n}}} \langle \xi \rangle^{10} |\widehat{\chi^2 A}(\tau, \xi)|^2 d\xi d\tau \\ &\leq (1 + R^{-\frac{3}{n}})^{10} \|\chi^2 A\|_{L^\infty(\mathbb{R}^{1+n})}^2 \left(\int_{-R}^R \int_{|\xi| \leq R^{-\frac{3}{n}}} d\xi d\tau \right) \end{aligned}$$

and hence

$$\int_{B_R \cap E_R} \langle \xi \rangle^{10} |\widehat{\chi^2 A}(\tau, \xi)|^2 d\xi d\tau \leq (2\pi)^{-(n+1)} 2^{11} T^2 |\Omega|^2 \|A\|_{L^\infty(Q)^n}^2 R^{-2}.$$

This and (4.9) yield

$$\int_{(\mathbb{R}^{1+n} \setminus B_R) \cup (B_R \cap E_R)} \langle \xi \rangle^{10} |\widehat{\chi^2 A}(\tau, \xi)|^2 d\xi d\tau \leq CR^{-2} \delta^{-2}. \quad (4.10)$$

On the other hand, putting $\epsilon := \|\Lambda_{A_1, q_1} - \Lambda_{A_2, q_2}\|$, we derive from (4.6) for all $(\tau, \xi) \in B_R \setminus E_R$, that

$$\begin{aligned} |\widehat{\chi^2 A}(\tau, \xi)|^2 &\leq C(R^{\frac{6}{n}+12} \delta^{-16} \sigma^{10} \epsilon^2 + R^{\frac{6}{n}+16} \delta^{-12} \sigma^{-2}) \\ &\leq C(R^{15} \delta^{-16} \sigma^{10} \epsilon^2 + R^{19} \delta^{-12} \sigma^{-2}) \end{aligned}$$

which involves

$$\int_{B_R \setminus E_R} \langle \xi \rangle^{10} |\widehat{\chi^2 A}(\tau, \xi)|^2 d\xi d\tau \leq C(R^{26+n} \delta^{-16} \sigma^{10} \epsilon^2 + R^{30+n} \delta^{-12} \sigma^{-2}).$$

It follows from this and (4.10) that

$$\|\chi^2 A\|_{L^2(0, T; H^5(\Omega))^n}^2 \leq C(R^{26+n} \delta^{-16} \sigma^{10} \epsilon^2 + R^{30+n} \delta^{-12} \sigma^{-2} + R^{-2} \delta^{-2}). \quad (4.11)$$

Further, by noticing that there exists a constant $C_0 > 0$ such that

$$\|A - \chi^2 A\|_{L^2(0, T; H^5(\Omega))^n} \leq C_0 |\Omega|^{1/2} \|1 - \chi^2\|_{L^2(0, T)} \|A\|_{W^{5, \infty}(Q)}$$

and taking advantage of the fact that the $[0, 1]$ -valued function $1 - \chi$ vanishes in $[2\delta, T - 2\delta]$, we get that $\|1 - \chi^2\|_{L^2(0, T)}^2 = \int_0^{2\delta} (1 - \chi^2(t))^2 dt + \int_{T-2\delta}^T (1 - \chi^2(t))^2 dt \leq 4\delta$. This entails

$$\|A - \chi^2 A\|_{L^2(0, T; H^5(\Omega))^n}^2 \leq 4C_0^2 |\Omega| \|A\|_{W^{5, \infty}(Q)}^2 \delta$$

and consequently

$$\|A\|_{L^2(0, T; H^5(\Omega))^n}^2 \leq C(\epsilon^2 R^{26+n} \delta^{-16} \sigma^{10} + R^{30+n} \delta^{-12} \sigma^{-2} + R^{-2} \delta^{-2} + \delta), \quad (4.12)$$

by invoking (4.11). Now, the strategy is to choose δ as a power of R so that $R^{-2} \delta^{-2} = \delta$, i.e. $\delta = R^{-\frac{2}{3}}$, and to do the same with σ , that is to take $\sigma = R^{\frac{16+3n}{6}}$, in such a way that the three last terms in the right hand side of (4.12) are equal to $R^{-\frac{2}{3}}$. Evidently, as we have $\delta \in (0, \frac{T}{4})$, by assumption, this requires that R be fixed in $(\frac{T}{4})^{-\frac{3}{2}}, \infty)$. Summing up, we infer from (4.12) that

$$\|A\|_{L^2(0, T; H^5(\Omega))^n}^2 \leq C(R^{230+6n} \epsilon^2 + R^{-\frac{2}{3}}). \quad (4.13)$$

Therefore, we get (1.7) with $r := \frac{1}{346+9n}$ for all $\epsilon \in (0, \epsilon_r)$, where $\epsilon_r := (\frac{T}{4})^{\frac{1}{2r}}$, upon choosing $R = \epsilon^{-3r}$ in (4.13), whereas $\|A\|_{L^2(0, T; H^5(\Omega))^n} \leq \frac{2M}{\epsilon_r} \epsilon^r$ for all $\epsilon \in [\epsilon_r, +\infty)$. This achieves the proof of (1.7).

5. PROOF OF THE STABILITY ESTIMATE (1.8)

Here we use the definitions and notations introduced in Sections 3 and 4, unless for the function β , which is no longer given by (3.10) but is rather defined by

$$\beta(t, x) := e^{-i(t\tau + x \cdot \xi)}, \quad (t, x) \in (0, T) \times \mathbb{R}^n.$$

As this definition formally coincides with (3.10) in the particular case where A is uniformly zero and $y = i\xi$, it is apparent that the estimates derived in Sections 3 and 4 remain valid with this specific choice of β .

Thus, in light of (3.11)–(3.16) and (4.3)–(4.4), it holds true that

$$\begin{aligned} &\left| \int_{\mathbb{R}^{1+n}} V u_{1,1} \overline{u_{2,1}}(t, x) dx dt \right| \\ &\leq C(\|A\|_{L^\infty(Q)^n} \langle \tau, \xi \rangle^8 \delta^{-6} \sigma + \|\Lambda_{A_2, q_2} - \Lambda_{A_1, q_1}\| \langle \tau, \xi \rangle^6 \delta^{-8} \sigma^6 + \langle \tau, \xi \rangle^6 \delta^{-4} \sigma^{-1}). \end{aligned} \quad (5.1)$$

Next, from the very definition of V , we have

$$\begin{aligned} \int_{\mathbb{R}^{1+n}} V u_{1,1} \overline{u_{2,1}}(t, x) dx dt &= \int_{\mathbb{R}^{1+n}} q u_{1,1} \overline{u_{2,1}}(t, x) dx dt - i \int_{\mathbb{R}^{1+n}} A \cdot \nabla(u_{1,1} \overline{u_{2,1}})(t, x) dx dt \\ &\quad - \int_{\mathbb{R}^{1+n}} A \cdot (A_1 + A_2) u_{1,1} \overline{u_{2,1}}(t, x) dx dt, \end{aligned}$$

by applying the Stokes formula. This, (3.11)–(3.16) and (5.1) yield

$$\left| \int_{\mathbb{R}^{1+n}} qu_{1,1}\overline{u_{2,1}}(t, x) dx dt \right| \quad (5.2)$$

$$\leq C \left(\|A\|_{L^\infty(Q)^n} \langle \tau, \xi \rangle^8 \delta^{-6} \sigma + \|\Lambda_{A_2, q_2} - \Lambda_{A_2, q_2}\| \langle \tau, \xi \rangle^6 \delta^{-8} \sigma^6 + \langle \tau, \xi \rangle^6 \delta^{-4} \sigma^{-1} \right). \quad (5.3)$$

On the other hand, we have

$$\int_{\mathbb{R}^{1+n}} qu_{11}\overline{u_{21}}(t, x) dx dt = \int_{\mathbb{R}^{1+n}} \chi^2(t) q(t, x) e^{-i(t\tau+x\cdot\xi)} e^{i \int_0^{+\infty} A(t, x+s\omega) \cdot \omega ds} dx dt,$$

whence

$$\begin{aligned} & \left| \int_{\mathbb{R}^{1+n}} \chi^2(t) q(t, x) e^{-i(t\tau+x\cdot\xi)} dx dt \right| \\ & \leq \left| \int_{\mathbb{R}^{1+n}} qu_{11}\overline{u_{21}}(t, x) dx dt \right| + \left| \int_{\mathbb{R}^{1+n}} q(t, x) e^{-i(t\tau+x\cdot\xi)} \left(e^{i \int_0^{+\infty} A(t, x+s\omega) \cdot \omega ds} - 1 \right) dx dt \right|. \end{aligned}$$

Thus, applying the mean value theorem, we get that

$$\left| \int_{\mathbb{R}^{1+n}} \chi^2(t) q(t, x) e^{-i(t\tau+x\cdot\xi)} dx dt \right| \leq \left| \int_{\mathbb{R}^{1+n}} qu_{11}\overline{u_{21}} dx dt \right| + C \|A\|_{L^\infty(Q)^n}.$$

Plugging (5.3) into the above estimate, we find that

$$\left| \widehat{\chi^2 q}(\tau, \xi) \right| \leq C \left(\|A\|_{L^\infty(Q)^n} \langle \tau, \xi \rangle^8 \delta^{-6} \sigma + \|\Lambda_{A_2, q_2} - \Lambda_{A_2, q_2}\| \langle \tau, \xi \rangle^6 \delta^{-8} \sigma^6 + \langle \tau, \xi \rangle^6 \delta^{-4} \sigma^{-1} \right). \quad (5.4)$$

The next step of the proof is to upper bound $\|A\|_{L^\infty(Q)^n}$ in terms of $\|\Lambda_{A_1, q_1} - \Lambda_{A_2, q_2}\|$. To this end, we pick $p \in (n+1, +\infty)$, apply the Sobolev embedding theorem (see e.g. [14, Corollary IX.14]) and find that $\|A\|_{L^\infty(Q)^3} \leq C \|A\|_{W^{1,p}(Q)^n}$. Interpolating, we obtain

$$\|A\|_{L^\infty(Q)^n} \leq C \|A\|_{W^{2,p}(Q)^n}^{1/2} \|A\|_{L^p(Q)^n}^{1/2} \leq C \|A\|_{L^p(Q)^n}^{1/2} \leq C \|A\|_{L^2(Q)^n}^{1/p}$$

and hence

$$\|A\|_{L^\infty(Q)^n} \leq C \|\Lambda_{A_1, q_1} - \Lambda_{A_2, q_2}\|^{r/p},$$

with the help of (1.7). Inserting the above estimate into (5.4) then yields

$$\left| \widehat{\chi^2 q}(\tau, \xi) \right| \leq C \left(\|\Lambda_{A_1, q_1} - \Lambda_{A_2, q_2}\|^{r/p} \langle \tau, \xi \rangle^8 \delta^{-6} \sigma + \|\Lambda_{A_2, q_2} - \Lambda_{A_2, q_2}\| \langle \tau, \xi \rangle^6 \delta^{-8} + \langle \tau, \xi \rangle^6 \delta^{-4} \sigma^{-1} \right)$$

and (1.8) follows from this by arguing in the same way as in the derivation of (1.7) from (4.12).

REFERENCES

- [1] G. ALESSANDRINI, *Stable determination of conductivity by boundary measurements*, Appl. Anal., **27** (1-3) (1988), 153-172.
- [2] C. BARDOS, G. LEBEAU AND J. RAUCH, *Sharp sufficient conditions for the observation, control and stabilization from the boundary*, SIAM J. Control Optim., **30** (1992), 1024-1165.
- [3] L. BAUDOIN AND J.-P. PUEL, *Uniqueness and stability in an inverse problem for the Schrödinger equation*, Inverse Problems, **18** (2002), 1537-1554.
- [4] M. BELLASSOUED, *Stable determination of coefficients in the dynamical Schrödinger equation in a magnetic field*, Inverse Problems, **33** (2017), 055009.
- [5] M. BELLASSOUED AND I. BEN AÏCHA, *Stable determination outside a cloaking region of two time-dependent coefficients in an hyperbolic equation from Dirichlet to Neumann map*, Jour. Math. Anal. Appl., **449** (2017), 46-76.
- [6] M. BELLASSOUED AND M. CHOULLI, *Logarithmic stability in the dynamical inverse problem for the Schrodinger equation by an arbitrary boundary observation*, J. Math. Pures Appl., **91** (2009), no. 3, 233-255.
- [7] M. BELLASSOUED AND M. CHOULLI, *Stability estimate for an inverse problem for the magnetic Schrödinger equation from the Dirichlet-to-Neumann map*, J. Funct. Anal., **258** (2010), 161-195.
- [8] M. BELLASSOUED AND D. DOS SANTOS FERREIRA, *Stable determination of coefficients in the dynamical anisotropic Schrödinger equation from the Dirichlet-to-Neumann map*, Inverse Problems, **26** (2010), 125010.
- [9] M. BELLASSOUED, Y. KIAN AND E. SOCCORSI, *An inverse stability result for non compactly supported potentials by one arbitrary lateral Neumann observation*, J. Diff. Equat., **260** (2016), no. 10, 7535-7562.
- [10] M. BELLASSOUED, Y. KIAN AND E. SOCCORSI, *An inverse problem for the magnetic Schrödinger equation in infinite cylindrical domains*, preprint arXiv:1605.06599.
- [11] I. BEN AÏCHA, *Stability estimate for hyperbolic inverse problem with time-dependent coefficient*, Inverse Problems, **31** (2015), 125010.

- [12] I. BEN AÏCHA, *Stability estimate for an inverse problem for the Schrödinger equation in a magnetic field with time-dependent coefficient*, preprint, arXiv:1608.00449.
- [13] I. BEN AÏCHA AND Y. MEJRI, *Simultaneous determination of the magnetic field and the electric potential in the Schrödinger equation by a finite number of boundary observations*, preprint, hal-01351217.
- [14] H. BREZIS, *Analyse Fonctionnelle. Théorie et applications*, Collection mathématiques appliquées pour la maîtrise, Masson, Paris 1993.
- [15] A. L. BUKHGEIM AND M. V. KLIBANOV, *Global uniqueness of class of multidimensional inverse problems*, Soviet Math. Dokl., **24** (1981), 244-247.
- [16] M. CHOULLI AND Y. KIAN, *Stability of the determination of a time-dependent coefficient in parabolic equations*, MCRF, **3** (2) (2013), 143-160.
- [17] M. CHOULLI AND Y. KIAN, *Logarithmic stability in determining the time-dependent zero order coefficient in a parabolic equation from a partial Dirichlet-to-Neumann map. Application to the determination of a nonlinear term*, preprint, arXiv:1605.08672.
- [18] M. CHOULLI, Y. KIAN, E. SOCCORSI, *Stable determination of time-dependent scalar potential from boundary measurements in a periodic quantum waveguide*, SIAM J. Math. Anal., **47** (6) (2015), 4536-4558.
- [19] M. CRISTOFOL AND E. SOCCORSI, *Stability estimate in an inverse problem for non-autonomous magnetic Schrödinger equations*, Applicable Analysis, **90** (2011), no. 10, 1499-1520.
- [20] D. DOS SANTOS FERREIRA, C. E. KENIG, J. SJÖSTRAND AND G. UHLMANN, *Determining a magnetic Schrödinger operator from partial Cauchy data*, Comm. Math. Phys., **271** (2007), no. 2, 467-488.
- [21] G. ESKIN, *Inverse problems for the Schrödinger equations with time-dependent electromagnetic potentials and the Aharonov-Bohm effect*, J. Math. Phys., **49** (2008), 022105.
- [22] G. ESKIN, *Inverse hyperbolic problems with time-dependent coefficients*, Commun. Partial Diff. Eqns., **32** (11) (2007), 1737-1758.
- [23] G. ESKIN, *Inverse problems for general second order hyperbolic equations with time-dependent coefficients*, preprint, arXiv:1503.00825.
- [24] G. ESKIN AND J. RALSTON, *Inverse scattering problem for the Schrödinger equation with magnetic potential at a fixed energy*, Commun. Math. Phys., **173** (1995), 199-224.
- [25] A. FAVINI, G. FAGNELLI AND R. M. MININNI, *New Prospects in Direct, Inverse and Control Problems for Evolution Equations*, Springer INdAM Series **10**.
- [26] K. FUJISHIRO AND Y. KIAN, *Determination of time dependent factors of coefficients in fractional diffusion equations*, MCRF, **6** (2016), 251-269.
- [27] P. GAITAN AND Y. KIAN, *A stability result for a time-dependent potential in a cylindrical domain*, Inverse Problems, **29** (6) (2013), 065006.
- [28] V. ISAKOV, *Completeness of products of solutions and some inverse problems for PDE*, J. Diff. Equat., **92** (1991), 305-316.
- [29] V. ISAKOV, *On uniqueness in inverse problems for semilinear parabolic equations*, Arch. Rat. Mech. Anal., **124** (1993), 1-12.
- [30] Y. KIAN, *Unique determination of a time-dependent potential for wave equations from partial data*, to appear in Annales de l'IHP (C) nonlinear analysis, doi:10.1016/j.anihpc.2016.07.003.
- [31] Y. KIAN, *Stability in the determination of a time-dependent coefficient for wave equations from partial data*, Jour. Math. Anal. Appl., **436** (2016), 408-428.
- [32] Y. KIAN, *A multidimensional Borg-Levinson theorem for magnetic Schrödinger operators with partial spectral data*, to appear in Journal of Spectral Theory, arXiv:1504.04514.
- [33] Y. KIAN, *Recovery of time-dependent damping coefficients and potentials appearing in wave equations from partial data*, SIAM J. Math. Anal., **48** (6) (2016), 4021-4046.
- [34] Y. KIAN AND L. OKSANEN, *Recovery of time-dependent coefficient on Riemannian manifold for hyperbolic equations*, preprint, arXiv:1606.07243.
- [35] Y. KIAN, Q. S. PHAN AND E. SOCCORSI, *Carleman estimate for infinite cylindrical quantum domains and application to inverse problems*, Inverse Problems, **30** (2014), 055016.
- [36] Y. KIAN, Q. S. PHAN AND E. SOCCORSI, *Hölder stable determination of a quantum scalar potential in unbounded cylindrical domains*, Jour. Math. Anal. Appl., **426** (2015), no. 1, 194-210.
- [37] J. L. LIONS AND E. MAGENES, *Problèmes aux limites non homogènes et applications*, Vol. I, Dunod, Paris, 1968.
- [38] J. L. LIONS AND E. MAGENES, *Problèmes aux limites non homogènes et applications*, Vol. II, Dunod, Paris, 1968.
- [39] N. MANDACHE, *Exponential instability in an inverse problem for the Schrödinger equation*, Inverse Problems, **17** (5) (2001), 1435-1444.
- [40] RAKESH AND A. G. RAMM, *Property C and an Inverse Problem for a Hyperbolic Equation*, J. Math. Anal. Appl., **156** (1991), 209-219.
- [41] A. G. RAMM AND J. SJÖSTRAND, *An inverse problem of the wave equation*, Math. Z., **206** (1991), 119-130.
- [42] P. STEFANOV, *Uniqueness of the multi-dimensional inverse scattering problem for time dependent potentials*, Math. Z., **201** (4) (1989), 541-559.
- [43] E. M. STEIN, *Singular integrals and differentiability properties of functions*, Princeton University Press, 1970.
- [44] L. TZOU, *Stability Estimate for the coefficients of magnetic Schrödinger equation from full and partial boundary measurements*, Commun. PDE, **11** (2008), 1911-1952.