

Carleman estimate for infinite cylindrical quantum domains and application to inverse problems

¹YAVAR KIAN, ²QUANG SANG PHAN, ³ERIC SOCCORSI

Abstract

We consider the inverse problem of determining the time independent scalar potential q of the dynamic Schrödinger equation in an infinite cylindrical domain Ω , from one Neumann boundary observation of the solution. Assuming that q is known outside some fixed compact subset of Ω , we prove that q may be Lipschitz stably retrieved by choosing the Dirichlet boundary condition of the system suitably. Since the proof is by means of a global Carleman estimate designed specifically for the Schrödinger operator acting in an unbounded cylindrical domain, the Neumann data is measured on an infinitely extended subboundary of the cylinder.

AMS 2010 Mathematics Subject Classification: 35R30.

Keywords: Inverse problem, Schrödinger equation, scalar potential, Carleman estimate, infinite cylindrical domain.

Contents

1	Introduction	2
1.1	What we are aiming for	2
1.2	Physical motivation: estimating the electrostatic quantum disorder in nanotubes	3
1.3	Existing papers	4
1.4	Main results	5
1.5	Outline	8
2	Analysis of the direct problem	8
2.1	Existence and uniqueness results	8
2.2	Linearized problem	9
2.3	Time symmetrization	10

¹Aix-Marseille Université, CNRS, CPT UMR 7332, 13288 Marseille, France & Université du Sud-Toulon-Var, 83957 La Garde, France. E-mail: yavar.kian@univ-amu.fr.

²Institute of Mathematics, Faculty of Mathematics and Computer Science, Jagiellonian University, 30-348 Kraków, Poland. E-mail: Quang.Sang.Phan@im.uj.edu.pl.

³Aix-Marseille Université, CNRS, CPT UMR 7332, 13288 Marseille, France & Université du Sud-Toulon-Var, 83957 La Garde, France. E-mail: eric.soccorsi@univ-amu.fr.

3	Carleman estimate for the dynamic Schrödinger operator	10
3.1	What is known in a bounded domain	11
3.2	Extension to the case of unbounded cylindrical domains	12
4	Proof of Theorem 1.1	15
4.1	Proof of the stability inequality (1.7)	15
4.1.1	Two auxiliary results	15
4.1.2	Completion of the proof	17
4.2	Proof of the stability estimate (1.8)	18

1 Introduction

1.1 What we are aiming for

In the present paper we consider the infinite cylindrical domain $\Omega = \omega \times \mathbb{R}$, where ω is a connected bounded open subset of \mathbb{R}^{n-1} , $n \geq 2$, with C^2 -boundary $\partial\omega$. Given $T > 0$ we examine the following initial boundary value problem

$$\begin{cases} -iu' - \Delta u + q(x)u = 0, & \text{in } Q := (0, T) \times \Omega, \\ u(0, x) = u_0(x), & x \in \Omega, \\ u(t, x) = g(t, x), & (t, x) \in \Sigma := (0, T) \times \Gamma, \end{cases} \quad (1.1)$$

where $\Gamma := \partial\omega \times \mathbb{R}$ and the ' stands for $\frac{\partial}{\partial t}$. Here u_0 (resp., g) is the initial (resp., boundary) condition associated to (1.1) and q is a function of $x \in \Omega$ only.

Since Γ is unbounded we make the boundary condition in the last line of (1.1) more precise. Writing $x := (x', x_n)$ with $x' := (x_1, \dots, x_{n-1}) \in \omega$ for every $x \in \Omega$ we extend the mapping

$$\begin{aligned} C_0^\infty((0, T) \times \mathbb{R}; H^2(\omega)) &\longrightarrow L^2((0, T) \times \mathbb{R}; H^{3/2}(\partial\omega)) \\ v &\mapsto [(t, x_n) \in (0, T) \times \mathbb{R} \mapsto v(t, \cdot, x_n)|_{\partial\omega}], \end{aligned} \quad (1.2)$$

to a bounded operator from $L^2((0, T) \times \mathbb{R}; H^2(\omega))$ into $L^2((0, T) \times \mathbb{R}; H^{3/2}(\partial\omega))$, denoted by γ_0 . Then for every $u \in C^0([0, T]; H^2(\Omega))$ the above mentioned boundary condition reads $\gamma_0 u = g$.

The main purpose of this article is to extend [3][Theorem 1], obtained for the dynamic Schrödinger operator in a bounded domain, to the case of the unbounded waveguide Ω . That is to say, to prove Lipschitz stability in the determination of the scalar potential q (assumed to be known outside some fixed compact subset of Ω) from one boundary measurement of the normal derivative of the solution u to (1.1).

The method of derivation of [3][Theorem 1] is by means of a Carleman estimate for the Schrödinger equation, which was established in [3][Proposition] for a bounded domain. Since the unknown part of q is compactly supported, it seems at first sight

quite reasonable to guess that this question (arising from the problem of evaluating the electrostatic quantum disorder in nanotubes, see subsection 1.2 below) could well be answered by adapting the above technique to some suitable truncation (for some bounded domain of \mathbb{R}^n) of u . Nevertheless we shall prove that such a strategy necessarily adds unexpected “control” terms (i.e. “volume observations” of the solution) in the right hand side of the corresponding stability inequality, and is therefore inaccurate.

It turns out that this inconvenience can be avoided upon substituting some specifically designed Carleman estimate for the Schrödinger equation in a unbounded cylindrical domain to the one of [3][Proposition 3]. This new global Carleman inequality, established in Proposition 3.3, is the main novelty of this paper as it is the main tool for generalizing the Lipschitz stability inequality of [3][Theorem 1] to the unbounded domain Ω under consideration.

It is worth noticing that the field of applications of Proposition 3.3 goes beyond the study of inverse compactly supported quantum scalar coefficients PDE problems examined in this framework. As a matter of facts we prove with the aid of Proposition 3.3 in a companion article [22], that scalar potentials which are not necessarily known outside some given compact subset of Ω , may nevertheless be Hölder stably retrieved from one boundary measurement of the normal derivative of u' .

Finally, let us mention that the Lipschitz stability estimate [3][Theorem 1] was established under the additional technical assumption that both u and u' are time square integrable bounded functions of the space variables. As we aim to generalize this result to the case of the unbounded domain Ω , we do the same here by supposing (1.6). Nevertheless we stress out that sufficient conditions on q and the initial state u_0 (which are not detailed in this text in order to avoid the inadequate expense of the size of the paper) ensuring assumption (1.6), can be found in [22][Theorem 1.1].

1.2 Physical motivation: estimating the electrostatic quantum disorder in nanotubes

The equations of (1.1) describe the evolution of the wave function of a charged particle (in a “natural” system of units where the various physical constants such as the mass and the electric charge of the particle are taken equal to one), under the influence of the “electric” potential q . Moreover the quantum motion of this particle is constrained by the waveguide Ω .

Carbon nanotubes exhibit unusual physical properties, which are valuable for electronics, optics and other fields of materials science and technology. These peculiar nanostructures have a length-to-diameter ratio up to $10^8/1$. This justifies why they may be rightfully modelled by infinite cylindrical domains such as Ω . Unfortunately their physical properties are commonly affected by the inevitable presence of electrostatic quantum disorder, see e. g. [11, 20]. This motivates for a closer look into the inverse problem of estimating the strength of the “electric impurity potential” q from the (partial) knowledge of the wave function u .

1.3 Existing papers

There is a wide mathematical literature dealing with uniqueness and stability in inverse coefficient problems related to partial differential equations, see e. g. [4, 7, 14, 18, 17, 23]. For the stationary (elliptic) Schrödinger equation, Bukhgeim and Uhlmann proved in [9] that the knowledge of the Dirichlet to Neumann (DN) map measured on some part of the boundary determines uniquely the potential. In dimension $n \geq 3$ this result was improved by Kenig, Sjöstrand and Uhlmann in [21] and L. Tzou proved in [27] that the electric potential (together with the magnetic field) depends stably on the Cauchy data even when the boundary measurement is taken only on a subset that is slightly larger than half of the boundary. Their method is essentially based on the construction of a rich set of “optics geometric solutions” to the Dirichlet problem. In these three papers, the knowledge of the DN map, taking infinitely many boundary measurements of the solution to the Schrödinger equation, is actually required.

The problem of stability in determining the time-independent electric potential in the dynamic Schrödinger equation from a single boundary measurement was treated by Baudouin and Puel in [3]. This result was improved by Mercado, Osses and Rosier in [26]. In these two papers, the main assumption is that the part of the boundary where the measurement is made satisfies a geometric condition related to geometric optics condition insuring observability (see Bardos, Lebeau and Rauch [2]). This geometric condition was relaxed in [5] under the assumption that the potential is known near the boundary.

In all the above mentioned articles the Schrödinger equation is defined in a bounded domain. In the present paper we rather investigate the problem of determining the scalar potential of the Schrödinger equation in an infinite cylindrical domain. There are only a few mathematical papers dealing with inverse boundary value problems in an unbounded domain available in the mathematical literature. In [25] Li and Uhlmann prove uniqueness in the determination of the scalar potential in an infinite slab from partial DN map. In [10] Cardoulis, Cristofol and Gaitan obtain Lipschitz stability from a single lateral measurement performed on one side of an unbounded strip.

More specifically for an inverse boundary value problem stated in the waveguide geometry examined in this work we refer to [12, 13] where stability is claimed for various coefficients of the Schrödinger equation from the knowledge of the DN map. Here we investigate the same type of problems, but in absence of any information given by the DN map, by a method based on an appropriate Carleman estimate. We refer to [1, 3, 27] for actual examples of this type inequalities designed for the Schrödinger equation. The original idea of using a Carleman estimate to solve inverse problems goes back to the pioneering article [8] by Bukhgeim and Klivanov. Since then this technique has then been widely and successfully used by numerous authors, see e.g. [3, 4, 6, 7, 12, 15, 16, 19, 23, 26], in the study of inverse wave propagation, elasticity or parabolic problems. The most recent survey about this method can be found in [24].

1.4 Main results

In this section we state the main results of this article and briefly comment on them.

For $M > 0$ and $p \in W^{2,\infty}(\Omega)$ fixed, we define the set of “admissible scalar potentials” as

$$\mathcal{Q}_M(p) := \{q \in W^{2,\infty}(\Omega; \mathbb{R}), \|q\|_{W^{2,\infty}(\Omega)} \leq M \text{ and } q(x) = p(x) \text{ for a.e. } x \in \Gamma\}.$$

Here the identity $q_\Gamma = p_\Gamma$ is understood in the sense of the usual trace operator from $L^\infty(\mathbb{R}, H^2(\omega))$ into $L^\infty(\mathbb{R}, H^{3/2}(\partial\omega))$. By selecting $q \in \mathcal{Q}_M(p)$ we thus enjoin fixed value to q on the boundary, which is the measurement on Γ of the scalar potential we want to determine. Similar (or stronger) “compatibility conditions” imposed on inverse problems coefficients have already been used in various contexts, see e.g. [5, 6, 14].

We are concerned with the stability issue around any $q_1 \in \mathcal{Q}_M(p)$, i.e. we want to upper bound the L^2 norm of $q_1 - q_2$ by some increasing function of the difference $u_1 - u_2$. That is to say that $q_2 \in \mathcal{Q}_M(p)$ and the solution u_j , for $j = 1, 2$, to (1.1), where q_j is substituted for q , are known, while q_1 is unknown.

Theorem 1.1. *For $p \in W^{2,\infty}(\Omega)$, $M > 0$, $\ell > 0$ and $\alpha > 0$ fixed, let $u_0 \in H^4(\Omega; \mathbb{R})$ obey*

$$u_0(x) \geq \alpha > 0, \quad x \in \omega \times (-\ell, \ell), \quad (1.3)$$

let $q_j \in \mathcal{Q}_M(p)$, $j = 1, 2$, fulfill

$$q_1(x) = q_2(x), \quad x \in \omega \times (\mathbb{R} \setminus (-\ell, \ell)), \quad (1.4)$$

and let u_j denote the $C^1([0, T]; H^2(\Omega) \cap H_0^1(\Omega)) \cap C^2([0, T]; L^2(\Omega))$ -solution to (1.1) associated to u_0 , $g = \gamma_0 G$ and q_j , where

$$G(t, x) := u_0(x) + it(\Delta - p)u_0(x), \quad (t, x) \in Q. \quad (1.5)$$

Assume that

$$u_2 \in H^1(0, T; L^\infty(\Omega)). \quad (1.6)$$

Then for every $L > \ell$, there exist $\Gamma_* \subset \partial\omega \times (-L, L)$ and a constant $C > 0$ depending only on L, T, M, ω and Γ_* , such that we have

$$\|q_1 - q_2\|_{L^2(\Omega)} \leq C \left(\|\partial_\nu(u'_1 - u'_2)\|_{L^2((0, T) \times \Gamma_*)} + \|u_1 - u_2\|_{H^1(0, T; H^1(\omega \times \mathcal{S}_L))} \right), \quad (1.7)$$

with $\mathcal{S}_L := (-L, -\ell) \cup (\ell, L)$.

Moreover there is a subboundary $\gamma_* \subset \partial\omega$, such that the estimate

$$\|q_1 - q_2\|_{L^2(\Omega)} \leq C' \|\partial_\nu u'_1 - \partial_\nu u'_2\|_{L^2((0, T) \times \gamma_* \times \mathbb{R})}, \quad (1.8)$$

holds for some positive constant C' depending only on ℓ, T, M, ω and γ_* .

Under the prescribed conditions (1.3)–(1.5) and assumption (1.6), the first statement (1.7) of Theorem 1.1 claims Lipschitz stability in the determination of the scalar potential appearing in the dynamic Schrödinger equation in Ω from two different observations of the solution u to (1.1).

- The first one is a lateral measurement on some subboundary of $\partial\omega \times (-L, L)$ of the normal derivative

$$\partial_\nu u(t, x) := \nabla u(t, x) \cdot \nu(x), \quad (t, x) \in \Sigma,$$

where ν is the outward unit normal to Γ . Since

$$\nu(x) = \nu(x') = \begin{pmatrix} \nu'(x') \\ 0 \end{pmatrix}, \quad x = (x', x_n) \in \Gamma, \quad (1.9)$$

where $\nu'(x') \in \mathbb{R}^{n-1}$ denotes the outgoing normal vector to $\partial\omega$ computed at x' , we notice that

$$\partial_\nu u(t, x) = \partial_{\nu'} u(t, x) := \nabla_{x'} u(t, x) \cdot \nu'(x'), \quad (t, x) = (t, x', x_n) \in \Sigma, \quad (1.10)$$

where $\nabla_{x'}$ stands for the gradient operator w.r.t. $x' \in \omega$.

- The second observation is an internal measurement of u which is performed in each of the two “slices” $\mathcal{S}_L^- := \omega \times (-L, -\ell)$ and $\mathcal{S}_L^+ := \omega \times (\ell, L)$ of Ω . Although the \mathbb{R}^n -Lebesgue measure of \mathcal{S}_L^\pm can be made arbitrarily small by taking L sufficiently close to ℓ , this “volume observation” cannot be removed from the right hand side of (1.7) by taking L asymptotically close to ℓ since the prefactor (i.e. the constant C) tends to infinity as L goes to ℓ .

The occurrence of this internal measurement of u in (1.7) is due to the unbounded geometry of Ω . More precisely this is a direct consequence of the technique used for the derivation of the stability inequality (1.7), which is by means of a global Carleman estimate for the Schrödinger equation in a bounded domain. Indeed, this strategy requires a cut off function with first derivative supported in $(-L, -\ell) \cup (\ell, L)$, which gives rise to the measurement of u in \mathcal{S}_L^\pm . Notice that the use of a Carleman estimate known to be valid in a bounded domain of \mathbb{R}^n only, was made possible here since the difference $q_1 - q_2$ is compactly supported in \mathbb{R}^n . A fact that follows from assumption (1.4) expressing that the scalar potential to be retrieved is known outside some fixed bounded set.

One way to get rid of both volume observations simultaneously is to use a global Carleman estimate specifically designed for the unbounded quantum waveguide Ω , which is stated in Proposition 3.3. This yields (1.8), implying that the electrostatic quantum potential is now Lipschitz stably retrieved in Ω from only one lateral measurement of the normal derivative $\partial_\nu u$ on some subboundary of Σ . This result is similar to the one obtained in a bounded domain (under the same assumption as (1.6)) by Baudouin and Puel, see [3][Theorem 1]. It should nevertheless be noticed that, contrarily to (1.7), and

despite of the fact that the scalar potential under identification is assumed to be known outside a compact set, the Neumann data required in the right hand side of (1.8) is measured on an infinitely extended subboundary of Σ .

Finally, it is worth mentioning that Theorem 1.1 applies for a wide class of subboundaries Γ_* . Indeed, given a C^2 -domain \mathcal{O} in \mathbb{R}^n obeying

$$\omega \times (-\ell, \ell) \subset \mathcal{O} \subset \omega \times (-L, L), \quad (1.11)$$

then this is the case for any $\Gamma_* \supset \{x \in \partial\mathcal{O}, (x - x_0) \cdot \nu_{\mathcal{O}}(x) \geq 0\}$, where $\nu_{\mathcal{O}}$ is the outward unit normal to $\partial\mathcal{O}$ and x_0 is arbitrary in $\mathbb{R}^n \setminus \overline{\mathcal{O}}$. The same remark holds true for $\gamma_* \supset \{x' \in \partial\omega, (x' - x'_0) \cdot \nu'(x') \geq 0\}$, where x'_0 is arbitrarily fixed in $\mathbb{R}^{n-1} \setminus \overline{\omega}$.

Theorem 1.1 immediately entails the following uniqueness result.

Corollary 1.2. *Under the conditions of Theorem 1.1, it holds true for all $q_j \in Q_M(p)$, $j = 1, 2$, that any of the two following assumptions*

$$\begin{cases} \partial_\nu u'_1(t, x) = \partial_\nu u'_2(t, x) & \text{for a.e. } (t, x) \in (0, T) \times \Gamma_* \\ u_1(t, x) = u_2(t, x) & \text{for a.e. } (t, x) \in (0, T) \times \omega \times \mathcal{S}_L, \end{cases} \quad (1.12)$$

or

$$\partial_\nu u'_1(t, x) = \partial_\nu u'_2(t, x) \text{ for a.e. } (t, x) \in (0, T) \times \gamma_* \times \mathbb{R}, \quad (1.13)$$

yields $q_1 = q_2$.

It turns out that Corollary 1.2 may actually be deduced from known results. Indeed, assuming (1.12), we have $u_1 = u_2$ on $(0, T) \times \partial\mathcal{O}$, where \mathcal{O} is the same as in (1.11), since u_1 coincides with u_2 on Σ , and

$$\partial_\nu u'_1(t, x) = \partial_\nu u'_2(t, x), \quad (t, x) \in (0, T) \times \tilde{\Gamma}, \quad (1.14)$$

for any $\tilde{\Gamma}$ obeying $\{x \in \partial\mathcal{O}, (x - x_0) \cdot \nu_{\mathcal{O}}(x) \geq 0\} \subset \tilde{\Gamma} \subset \Gamma_* \cup (\partial\mathcal{O} \cap \mathcal{S}_L)$. Hence the restriction of u_j , $j = 1, 2$, to $(0, T) \times \mathcal{O}$, is solution to the following boundary value problem

$$\begin{cases} -iu' - \Delta u_j + q_j(x)u_j = 0, & \text{in } (0, T) \times \mathcal{O}, \\ u_j(0, x) = u_0(x), & x \in \mathcal{O}, \\ u_j(t, x) = \tilde{g}(t, x), & (t, x) \in (0, T) \times \partial\mathcal{O}, \end{cases}$$

where \tilde{g} is a function that coincides with g on $(0, T) \times \partial\omega \times (-\ell, \ell)$. Thus, applying [3][Theorem 1], we deduce from (1.14) that $q_1(x) = q_2(x)$ for a.e. $x \in \mathcal{O}$, which entails $q_1 = q_2$ a.e. in Ω by (1.4).

Quite similarly we may derive uniqueness from (1.4) and (1.13) by means of the unique continuation [5][Lemma 2.1 of the solution to the Schrödinger equation. Moreover the same remark applies for the unique continuation result [29][Theorem 1.2] as well, that was obtained from a Carleman estimate for the Schrödinger equation with quadratic weights, which are neither used in [3, 5] nor in this paper.

Anyhow, we stress out that the main purpose of this article is the stability issue in the inverse problem of determining the scalar potential of the Schrödinger equation in an unbounded cylindrical domain, which, as far as we know, cannot be treated by existing results.

1.5 Outline

The paper is organized as follows. Section 2 deals with the direct problem associated to (1.1). Namely subsection 2.1 gathers existence and uniqueness results for the solution to the dynamic Schrödinger equation in the infinite domain Ω and subsection 2.2 is devoted to the study of the direct problem for the linearized system associated to (1.1). In subsection 2.3 the corresponding solution is suitably extended to a function of $[-T, T] \times \Omega$ which is continuous with respect to the time variable t . This is required by the method used in the proof of the stability inequalities (1.7) and (1.8), which is the purpose of Section 4. It is by means of the global Carleman estimate for the Schrödinger equation in a bounded domain (resp., in an unbounded waveguide) stated in Proposition 3.2 (resp., Proposition 3.3) for (1.7) (resp., for (1.8)). Finally section 4 contains the completion of the proof of (1.7)-(1.8).

2 Analysis of the direct problem

In this section we establish some existence, uniqueness and regularity properties of the solution to (1.1) needed for the analysis of the inverse problem in section 3.

2.1 Existence and uniqueness results

This subsection gathers two existence and uniqueness results for (1.1). We start by recalling from [13][Proposition 2.1] the following:

Proposition 2.1. *Let $M > 0$. Then for all $q \in \mathcal{Q}_M$, $v_0 \in \mathbf{H}_0^1(\Omega) \cap \mathbf{H}^2(\Omega)$ and $f \in W^{1,1}(0, T; \mathbf{L}^2(\Omega))$, there exists a unique solution $v \in Z_0 := C([0, T]; \mathbf{H}_0^1(\Omega) \cap \mathbf{H}^2(\Omega)) \cap C^1([0, T]; \mathbf{L}^2(\Omega))$ to the boundary value problem*

$$\begin{cases} -iv' - \Delta v + qv = f, & \text{in } Q, \\ v(0, x) = v_0, & x \in \Omega, \\ v(t, x) = 0, & (t, x) \in \Sigma. \end{cases}$$

Moreover we have

$$\|v\|_{Z_0} \leq C (\|v_0\|_{\mathbf{H}^2(\Omega)} + \|f\|_{W^{1,1}(0, T; \mathbf{L}^2(\Omega))}),$$

for some constant $C > 0$ depending only on ω , T and M .

Let the space

$$\mathcal{X}_0 := \gamma_0(W^{2,2}(0, T; \mathbf{H}^2(\Omega))), \tag{2.15}$$

be equipped with the following quotient norm

$$\|g\|_{\mathcal{X}_0} = \inf\{\|G\|_{W^{2,2}(0, T; \mathbf{H}^2(\Omega))}; G \in W^{2,2}(0, T; \mathbf{H}^2(\Omega)) \text{ satisfies } \gamma_0 G = g\}.$$

Evidently, each $g \in \mathcal{X}_0$ admits an extension $G_0 \in W^{2,2}(0, T; H^2(\Omega))$ verifying

$$\|G_0\|_{W^{2,2}(0, T; H^2(\Omega))} \leq 2\|g\|_{\mathcal{X}_0}.$$

Further, put

$$\mathcal{L} := \{(u_0, g) \in H^2(\Omega) \times \mathcal{X}_0; u_0 = g(0, \cdot) \text{ on } \Gamma\}. \quad (2.16)$$

Then, setting $u = v + G_0$, where v is defined by Proposition 2.1 for $f = i\partial_t G_0 + \Delta G_0 - qG_0$ and $v_0 = u_0 - G_0(0, \cdot)$, we obtain the following existence and uniqueness result, which is similar to [13][Corollary 2.1].

Theorem 2.2. *Let $M > 0$. Then for all $q \in \mathcal{Q}_M$ and all $(u_0, g) \in \mathcal{L}$ there exists a unique solution $u \in Z := C([0, T]; H^2(\Omega)) \cap C^1([0, T]; L^2(\Omega))$ to the boundary value problem (1.1). Moreover we have the following estimate*

$$\|u\|_Z \leq C(\|u_0\|_{H^2(\Omega)} + \|g\|_{\mathcal{X}_0}), \quad (2.17)$$

where C is some positive constant depending only on ω , T and M .

2.2 Linearized problem

In this subsection we establish some useful regularity properties of the solution to the linearized problem associated to (1.1) by applying the existence and uniqueness results stated in subsection 2.1.

To this purpose, we let p , M , q_j , $j = 1, 2$, u_0 , g and G be the same as in Theorem 1.1. Since $G \in W^{2,2}(0, T; H^2(\Omega))$ and $g(0, \cdot) = u_0$ on Γ , we have $(u_0, g) \in \mathcal{L}$. Applying Theorem 2.2 there is thus a unique solution $u_j \in Z$, $j = 1, 2$, to the following system

$$\begin{cases} -iu'_j - \Delta u_j + q_j(x)u_j = 0, & \text{in } Q, \\ u_j(0, x) = u_0(x), & x \in \Omega, \\ u_j(t, x) = g(t, x), & (t, x) \in \Sigma. \end{cases} \quad (2.18)$$

Further, differentiating (2.18) with respect to t for $j = 2$, we obtain that u'_2 is solution to the boundary value problem

$$\begin{cases} -iu''_2 - \Delta u'_2 + q_2(x)u'_2 = 0, & \text{in } Q, \\ u'_2(0, x) = \tilde{u}_0(x), & x \in \Omega, \\ u'_2(t, x) = g'(t, x), & (t, x) \in \Sigma, \end{cases} \quad (2.19)$$

where $\tilde{u}_0 := i(\Delta - q_2)u_0$. Since $G'(t, x) = i(\Delta - p)(x)$ for a.e. $(t, x) \in Q$ by (1.5) then we have $G' \in W^{2,2}(0, T; H^2(\Omega))$ and we get $g'(0, x) = \tilde{u}_0(x)$ for a.e. $x \in \Gamma$ from the identity $q_2 = p$ on Γ . Consequently $(\tilde{u}_0, g') \in \mathcal{L}$. Therefore $u'_2 \in Z$ from (2.19) and Theorem 2.2. As a consequence we have

$$u_2 \in C^1([0, T]; H^2(\Omega)) \cap C^2([0, T]; L^2(\Omega)).$$

Moreover, it follows from (2.18) that $u := u_1 - u_2$ is solution to the linearized system

$$\begin{cases} -iu' - \Delta u + q_1 u = f, & \text{in } Q, \\ u(0, x) = 0, & x \in \Omega, \\ u(t, x) = 0, & (t, x) \in \Sigma, \end{cases} \quad (2.20)$$

with $f := (q_2 - q_1)u_2$. Since $u_2 \in Z$ and $q_1 - q_2 \in L^\infty(\Omega)$ then $f \in W^{1,1}(0, T; L^2(\Omega))$ so we find that $u \in Z_0$ by Proposition 2.1.

Last, we deduce from (2.20) that $v := u'$ satisfies

$$\begin{cases} -iv' - \Delta v + q_1 v = f', & \text{in } Q, \\ v(0, x) = i(q_2 - q_1)(x)u_0(x), & x \in \Omega. \\ v(t, x) = 0, & (t, x) \in \Sigma, \end{cases} \quad (2.21)$$

with $f' = (q_2 - q_1)u_2' \in W^{1,1}(0, T; L^2(\Omega))$. Since $q_1 = q_2$ on Γ we get that $i(q_2 - q_1)u_0 \in H_0^1(\Omega) \cap H^2(\Omega)$. Therefore $v = u_1' - u_2' \in Z_0$ by Proposition 2.1. Bearing in mind that $u_2' \in Z$ we deduce from this that $u_1' \in Z$. Summing up, we have obtained that

$$u_j \in C^1([0, T]; H^2(\Omega)) \cap C^2([0, T]; L^2(\Omega)), \quad j = 1, 2.$$

2.3 Time symmetrization

This subsection is devoted to showing that the solution $v \in Z_0$ to (2.21) may be extended to a solution of the same system over the time span $[-T, T]$.

To this end we extend v (resp., f') on $(-T, 0) \times \Omega$ by setting $v(t, x) := \overline{-v(-t, x)}$ (resp., $f'(t, x) := -\overline{f'(-t, x)}$) for $(t, x) \in (-T, 0) \times \Omega$. Since the initial conditions $v(0, \cdot)$ and $f'(0, \cdot)$ are purely complex valued according to (2.21), the mappings $t \mapsto v(t, x)$ and $t \mapsto f'(t, x)$ are thus continuous at $t = 0$ for a.e. $x \in \Omega$. Therefore, we have $v \in C^0([-T, T]; H_0^1(\Omega) \cap H^2(\Omega)) \cap C^1([-T, T]; L^2(\Omega))$ and $f' \in W^{1,1}(-T, T; L^2(\Omega))$. Moreover v is solution to (2.21) over the whole time span $(-T, T)$:

$$\begin{cases} -iv' - \Delta v + q_1 v = f', & \text{in } Q_e := (-T, T) \times \Omega, \\ v(0, x) = i(q_2 - q_1)(x)u_0(x), & x \in \Omega. \\ v(t, x) = 0, & (t, x) \in \Sigma_e := (-T, T) \times \Gamma. \end{cases} \quad (2.22)$$

3 Carleman estimate for the dynamic Schrödinger operator

The main purpose of this section is to derive a global Carleman estimate for the Schrödinger operator in an unbounded cylindrical domain $\mathcal{O} \times \mathbb{R}$, where \mathcal{O} is a bounded domain of \mathbb{R}^m , $m \geq 1$, from the same type of results obtained for \mathcal{O} by Baudouin and Puel in [3][Proposition 3].

3.1 What is known in a bounded domain

Given an arbitrary bounded domain $\mathcal{O} \supset \mathbb{R}^m$ with C^2 boundary, we consider the Schrödinger operator acting in $(C_0^\infty)'((-T, T) \times \mathcal{O})$,

$$L_m := -i\partial_t - \Delta, \quad (3.23)$$

where Δ denotes the Laplacian in \mathcal{O} . Here we use the subscript m to emphasize the fact that the operator L_m acts in a subdomain of $(-T, T) \times \mathbb{R}^m$.

Next we introduce a function $\tilde{\beta} \in C^4(\overline{\mathcal{O}}; \mathbb{R}_+)$ and an open subset $\Gamma_{\mathcal{O}}$ of $\partial\mathcal{O}$, satisfying the following conditions:

Assumption 3.1.

- (i) $\exists C_0 > 0$ such that the estimate $|\nabla\tilde{\beta}(x)| \geq C_0$ holds for all $x \in \mathcal{O}$;
- (ii) $\partial_\nu\tilde{\beta}(x) := \nabla\tilde{\beta}(x) \cdot \nu(x) < 0$ for all $x \in \partial\mathcal{O} \setminus \Gamma_{\mathcal{O}}$, where ν is the outward unit normal vector to $\partial\mathcal{O}$;
- (iii) $\exists \Lambda_1 > 0, \exists \epsilon > 0$ such that we have $\lambda|\nabla\tilde{\beta}(x) \cdot \zeta|^2 + D^2\tilde{\beta}(x, \zeta, \zeta) \geq \epsilon|\zeta|^2$ for all $\zeta \in \mathbb{R}^m, x \in \mathcal{O}$ and $\lambda > \Lambda_1$, where $D^2\tilde{\beta}(x) := \left(\frac{\partial^2\tilde{\beta}(x)}{\partial x_i \partial x_j} \right)_{1 \leq i, j \leq m}$ and $D^2\tilde{\beta}(x, \zeta, \zeta)$ denotes the \mathbb{R}^m -scalar product of $D^2\tilde{\beta}(x)\zeta$ with ζ .

Notice that there are actual functions $\tilde{\beta}$ verifying Assumption 3.1, such as $\mathcal{O} \ni x \mapsto |x - x_0|^2$, where x_0 is arbitrary in $\mathbb{R}^m \setminus \overline{\mathcal{O}}$ and $\Gamma_{\mathcal{O}} \supset \{x \in \partial\mathcal{O}, (x - x_0) \cdot \nu(x) \geq 0\}$.

Further we put

$$\beta := \tilde{\beta} + K, \text{ where } K := r\|\tilde{\beta}\|_\infty \text{ for some } r > 1, \quad (3.24)$$

and define the two following weight functions for $\lambda > 0$:

$$\varphi(t, x) := \frac{e^{\lambda\beta(x)}}{(T+t)(T-t)} \text{ and } \eta(t, x) := \frac{e^{2\lambda K} - e^{\lambda\beta(x)}}{(T+t)(T-t)}, \quad (t, x) \in (-T, T) \times \mathcal{O}. \quad (3.25)$$

Finally, for all $s > 0$ we introduce the two following operators acting in $(C_0^\infty)'((-T, T) \times \mathcal{O})$:

$$M_{1,m} := i\partial_t + \Delta + s^2|\nabla\eta|^2 \text{ and } M_{2,m} := is\eta' + 2s\nabla\eta \cdot \nabla + s(\Delta\eta). \quad (3.26)$$

It is easily seen that $M_{1,m}$ (resp., $M_{2,m}$) is the adjoint (resp., skew-adjoint) part of the operator $e^{-s\eta}L_m e^{s\eta}$, where L_m is given by (3.23).

Having said that we may now state the following global Carleman estimate, borrowed from [3][Proposition 3], which is the main tool for the proof of Theorem 1.1.

Proposition 3.2. *Let β be given by (3.24), where $\tilde{\beta} \in C^4(\overline{\mathcal{O}}; \mathbb{R}_+)$ fulfills Assumption 3.1, let φ and η be as in (3.25), and let $L_m, M_{1,m}$ and $M_{2,m}$ be defined by (3.23)-(3.26).*

Then we may find $s_0 > 0$ and constant $C > 0$, depending only on T , \mathcal{O} and $\Gamma_{\mathcal{O}}$, such that the estimate

$$\begin{aligned} & s \|e^{-s\eta} \nabla w\|_{L^2((-T,T) \times \mathcal{O})}^2 + s^3 \|e^{-s\eta} w\|_{L^2((-T,T) \times \mathcal{O})}^2 + \sum_{j=1,2} \|M_{j,m} e^{-s\eta} w\|_{L^2((-T,T) \times \mathcal{O})}^2 \\ & \leq C \left(s \|e^{-s\eta} \varphi^{1/2} (\partial_\nu \beta)^{1/2} \partial_\nu w\|_{L^2((-T,T) \times \Gamma_{\mathcal{O}})}^2 + \|e^{-s\eta} L_m w\|_{L^2((-T,T) \times \mathcal{O})}^2 \right), \end{aligned} \quad (3.27)$$

holds for any real number $s \geq s_0$ and any function $w \in L^2(-T, T; H_0^1(\mathcal{O}))$ satisfying $L_m w \in L^2((-T, T) \times \mathcal{O})$ and $\partial_\nu w \in L^2(-T, T; L^2(\Gamma_{\mathcal{O}}))$.

The proof of Proposition 3.2 can be found in [3][section 2].

3.2 Extension to the case of unbounded cylindrical domains

Let ω be $\Omega = \omega \times \mathbb{R}$ be the same as in section 1. We use the same notation as in section 1 and section 2, i.e. every point $x \in \Omega$ is written $x = (x', x_n)$ with $x' = (x_1, \dots, x_{n-1}) \in \omega$, and we note $Q_\varepsilon = (-T, T) \times \Omega$. We shall now derive from Proposition 3.2 a global Carleman estimate for the Schrödinger operator

$$L_n := -i\partial_t - \Delta, \quad (3.28)$$

acting in $(C_0^\infty)'((-T, T) \times \Omega)$. Here $\Delta := \Delta_{x'} + \partial_{x_n}^2$ where $\Delta_{x'} := \sum_{j=1}^{n-1} \partial_{x_j}^2$ is the Laplacian in ω . Similarly, we denote by $\nabla_{x'}$ the gradient operator w.r.t. $x' \in \omega$.

Proposition 3.3. *For $n \geq 2$ fixed, let $\tilde{\beta}$ and $\gamma_* := \Gamma_{\mathcal{O}}$ obey Assumption 3.1 with $\mathcal{O} = \omega$ and $m = n - 1$, and let β be given by (3.24). Extend β to $\Omega = \omega \times \mathbb{R}$ by setting $\beta(x) = \beta(x')$ for all $x = (x', x_n) \in \Omega$. Define φ and η (resp., the operators $M_{j,n}$ for $j = 1, 2$) by (3.25) (resp., (3.26)), upon substituting Ω for \mathcal{O} and n for m . Then there are two constants $s_0 > 0$ and $C > 0$, depending only on T , ω and γ_* , such that the estimate*

$$\begin{aligned} & s \|e^{-s\eta} \nabla_{x'} v\|_{L^2(Q_\varepsilon)}^2 + s^3 \|e^{-s\eta} v\|_{L^2(Q_\varepsilon)}^2 + \sum_{j=1,2} \|M_{j,n} e^{-s\eta} v\|_{L^2(Q_\varepsilon)}^2 \\ & \leq C \left(s \|e^{-s\eta} \varphi^{1/2} (\partial_\nu \beta)^{1/2} \partial_\nu v\|_{L^2((-T,T) \times \gamma_* \times \mathbb{R})}^2 + \|e^{-s\eta} L_n v\|_{L^2(Q_\varepsilon)}^2 \right), \end{aligned} \quad (3.29)$$

holds for all $s \geq s_0$ and any function $v \in L^2(-T, T; H_0^1(\Omega))$ verifying $L_n v \in L^2(Q_\varepsilon)$ and $\partial_\nu v \in L^2(-T, T; L^2(\gamma_* \times \mathbb{R}))$.

Proof. Since the function β does not depend on the infinite variable x_n then the same is true for φ and η according to (3.25). Therefore we have

$$M_{1,n} = i\partial_t + \Delta + s^2 |\nabla_{x'} \eta|^2 \text{ and } M_{2,n} = is\eta' + 2s(\nabla_{x'} \eta) \cdot \nabla_{x'} + s(\Delta_{x'} \eta), \quad (3.30)$$

We denote by \mathcal{F}_{x_n} the partial Fourier transform with respect to x_n , i.e.

$$(\mathcal{F}_{x_n}\psi)(x', k) = \widehat{\psi}(x', k) := \frac{1}{(2\pi)^{1/2}} \int_{\mathbb{R}} e^{-ikx_n} \psi(x', x_n) dx_n, \quad \psi \in L^2(\Omega), \quad x' \in \omega, \quad k \in \mathbb{R},$$

and consider the unitary group $U_t = e^{-it\partial_{x_n}^2}$ in $L^2(\Omega)$, defined by

$$U_t := \mathcal{F}_{x_n}^{-1} e^{itk^2} \mathcal{F}_{x_n}, \quad t \in \mathbb{R},$$

where $\mathcal{F}_{x_n}^{-1}$ is the inverse transform to \mathcal{F}_{x_n} and e^{itk^2} stands for the multiplier by the corresponding complex number. Therefore we have

$$(U_t\psi)(x) = e^{-it\partial_{x_n}^2} \psi(x) := \mathcal{F}_{x_n}^{-1} \left(e^{itk^2} \widehat{\psi}(\cdot, k) \right) (x), \quad \psi \in L^2(\Omega), \quad x \in \Omega, \quad t \in \mathbb{R}. \quad (3.31)$$

We notice from (3.31) that

$$[i\partial_t, U_t] = \partial_{x_n}^2 U_t = U_t \partial_{x_n}^2, \quad t \in \mathbb{R}, \quad (3.32)$$

since $\partial_{x_n}^2$ commutes with U_t , and that

$$[\partial_{x_j}, U_t] = [e^{-s\eta}, U_t] = 0, \quad j = 1, \dots, n-1 \text{ and } s, t \in \mathbb{R}, \quad (3.33)$$

since η does not depend on x_n . Let us now introduce the function

$$w(t, x) := U_t v(t, x), \quad (t, x) \in Q_e. \quad (3.34)$$

Then we have $w(\cdot, x_n) \in L^2(-T, T; H_0^1(\omega))$ and $\partial_{\nu'} w(\cdot, x_n) \in L^2(-T, T; L^2(\gamma_*))$ for a.e. $x_n \in \mathbb{R}$. Further, upon differentiating (3.34) w.r.t. t , we deduce from (3.32) that

$$i\partial_t w = U_t (i\partial_t v + \partial_{x_n}^2 v). \quad (3.35)$$

This and the first commutator identity in (3.33) yields

$$(-i\partial_t - \Delta_{x'}) w(t, x) = U_t L_n v(t, x), \quad (t, x) \in Q_e,$$

which entails

$$L_{n-1} w(\cdot, x_n) = U_t L_n v(\cdot, x_n), \quad x_n \in \mathbb{R}. \quad (3.36)$$

As a consequence we have $L_{n-1} w(\cdot, x_n) \in L^2((-T, T) \times \omega)$ for a.e. $x_n \in \mathbb{R}$, whence

$$\begin{aligned} & s \|e^{-s\eta} \nabla_{x'} w(\cdot, x_n)\|_{L^2((-T, T) \times \omega)}^2 + s^3 \|e^{-s\eta} w(\cdot, x_n)\|_{L^2((-T, T) \times \omega)}^2 \\ & + \sum_{j=1,2} \|M_{j,n-1} e^{-s\eta} w(\cdot, x_n)\|_{L^2((-T, T) \times \omega)}^2 \\ & \leq C \left(s \|e^{-s\eta} \varphi^{1/2} (\partial_{\nu'} \beta)^{1/2} \partial_{\nu'} w(\cdot, x_n)\|_{L^2((-T, T) \times \gamma_*)}^2 + \|e^{-s\eta} L_{n-1} w(\cdot, x_n)\|_{L^2((-T, T) \times \omega)}^2 \right), \end{aligned} \quad (3.37)$$

for every $s \geq s_0$, by Proposition 3.2.

The next step of the proof involves noticing from (3.33)–(3.35) that

$$e^{-s\eta}w(\cdot, x_n) = U_t e^{-s\eta}v(\cdot, x_n), \quad e^{-s\eta}\nabla_{x'}w(\cdot, x_n) = U_t e^{-s\eta}\nabla_{x'}v(\cdot, x_n), \quad (3.38)$$

and, with reference to (3.30), that

$$M_{j,n-1}e^{-s\eta}w(\cdot, x_n) = U_t M_{j,n}e^{-s\eta}v(\cdot, x_n), \quad j = 1, 2, \quad (3.39)$$

for each $t \in (-T, T)$ and a.e. $x_n \in \mathbb{R}$. Similarly, we get

$$e^{-s\eta}L_{n-1}w(\cdot, x_n) = U_t e^{-s\eta}L_n v(\cdot, x_n), \quad t \in (-T, T), \quad x_n \in \mathbb{R}, \quad (3.40)$$

by combining the second identity of (3.33) with (3.36). Finally, bearing in mind that φ and β are independent of x_n , we deduce from (1.9)–(1.10) and the second part of (3.33) that

$$e^{-s\eta}\varphi^{1/2}(\partial_\nu\beta)^{1/2}\partial_\nu w(\cdot, x_n) = U_t e^{-s\eta}\varphi^{1/2}(\partial_\nu\beta)^{1/2}\partial_\nu v(\cdot, x_n), \quad t \in (-T, T), \quad x_n \in \mathbb{R}. \quad (3.41)$$

Therefore, putting (3.37)–(3.41) together, we obtain for $s \geq s_0$ and a.e. $x_n \in \mathbb{R}$ that

$$\begin{aligned} & s\|U_t e^{-s\eta}\nabla_{x'}v(\cdot, x_n)\|_{L^2((-T,T)\times\omega)}^2 + s^3\|U_t e^{-s\eta}v(\cdot, x_n)\|_{L^2((-T,T)\times\omega)}^2 \\ & + \sum_{j=1,2} \|U_t M_{j,n}e^{-s\eta}v(\cdot, x_n)\|_{L^2((-T,T)\times\omega)}^2 \\ & \leq C \left(s\|U_t e^{-s\eta}\varphi^{1/2}(\partial_\nu\beta)^{1/2}\partial_\nu v(\cdot, x_n)\|_{L^2((-T,T)\times\gamma_*)}^2 + \|U_t e^{-s\eta}L_n v(\cdot, x_n)\|_{L^2((-T,T)\times\omega)}^2 \right). \end{aligned}$$

Integrating the above inequality w.r.t. x_n over \mathbb{R} and taking into account that the operator U_t is isometric in $L^2(\Omega)$ for each $t \in (-T, T)$, we find that

$$\begin{aligned} & s\|e^{-s\eta}\nabla_{x'}v\|_{L^2(Q_e)}^2 + s^3\|e^{-s\eta}v\|_{L^2(Q_e)}^2 + \sum_{j=1,2} \|M_{j,n}e^{-s\eta}v\|_{L^2(Q_e)}^2 \\ & \leq C \left(s\|U_t e^{-s\eta}\varphi^{1/2}(\partial_\nu\beta)^{1/2}\partial_\nu v\|_{L^2((-T,T)\times\gamma_*\times\mathbb{R})}^2 + \|e^{-s\eta}L_n v\|_{L^2(Q_e)}^2 \right), \quad (3.42) \end{aligned}$$

provided $s \geq s_0$. Last, setting $\Phi := e^{-s\eta}\varphi^{1/2}(\partial_\nu\beta)^{1/2}\partial_\nu v$ and using the unitarity of the operator \mathcal{F}_{x_n} in $L^2(\mathbb{R})$, we derive from (3.31) that

$$\begin{aligned} & \|U_t\Phi\|_{L^2((-T,T)\times\gamma_*\times\mathbb{R})}^2 = \int_{-T}^T \int_{\gamma_*} \|\mathcal{F}_{x_n}^{-1}e^{itk}\widehat{\Phi}(t, x', \cdot)\|_{L^2(\mathbb{R})}^2 dx' dt \\ & = \int_{-T}^T \int_{\gamma_*} \|e^{itk}\widehat{\Phi}(t, x', \cdot)\|_{L^2(\mathbb{R})}^2 dx' dt = \int_{-T}^T \int_{\gamma_*} \|\widehat{\Phi}(t, x', \cdot)\|_{L^2(\mathbb{R})}^2 dx' dt \\ & = \|\Phi\|_{L^2((-T,T)\times\gamma_*\times\mathbb{R})}^2, \end{aligned}$$

so (3.29) follows immediately from this and (3.42). \square

4 Proof of Theorem 1.1

In view of section 3 we are now in position to solve the inverse problem under study. More precisely we apply the Bugkhgeim-Klibanov method presented in [8] to prove the stability inequality (1.7) (resp., the estimate (1.8)) with the aid of Proposition 3.2 (resp., Proposition 3.3).

4.1 Proof of the stability inequality (1.7)

We first introduce the following notation used throughout the entire section: for all $d > 0$ we write $\Omega_d := \omega \times (-d, d)$.

Fix $L > \ell$, set $r := (L + \ell)/2$ and consider a domain \mathcal{O} in \mathbb{R}^n , with C^2 boundary $\partial\mathcal{O}$, obeying

$$\Omega_\ell \subset \Omega_r \subset \mathcal{O} \subset \Omega_L. \quad (4.43)$$

We next choose $\chi \in C_0^\infty(\mathbb{R}_{x_n}, [0, 1])$ such that

$$\chi(x_n) := \begin{cases} 1 & \text{if } |x_n| \leq \ell \\ 0 & \text{if } |x_n| \geq r. \end{cases}$$

Put $w := \chi v$, where v is the $C^0([-T, T]; H_0^1(\Omega) \cap H^2(\Omega)) \cap C^1([-T, T]; L^2(\Omega))$ -solution to (2.22); Then the restriction $w|_{\mathcal{O}}$ of w to \mathcal{O} satisfies

$$w|_{\mathcal{O}} \in C^0([-T, T]; H_0^1(\mathcal{O}) \cap H^2(\mathcal{O})) \cap C^1([-T, T]; L^2(\mathcal{O})) \quad (4.44)$$

and is solution to the following system

$$\begin{cases} -iw' - \Delta w + q_1 w = \chi f' - K v, & \text{in } (-T, T) \times \mathcal{O}, \\ w(0, x) = i\chi(q_2 - q_1)(x)u_0(x), & x \in \mathcal{O}. \\ w(t, x) = 0, & (t, x) \in (-T, T) \times \partial\mathcal{O}, \end{cases} \quad (4.45)$$

where $K := [\Delta, \chi] = \ddot{\chi} + 2\dot{\chi}\partial_{x_n}$. Here $\dot{\chi}$ (resp., $\ddot{\chi}$) is a shorthand for the first (resp., second) derivative of χ .

As a preamble to the proof of the stability inequality (1.7) we first establish two elementary technical results.

4.1.1 Two auxiliary results

Lemma 4.1. *For $s > 0$, let $\phi := e^{-s\eta} w$, where w is defined by (4.44)-(4.45). Then we have*

$$J := \|e^{-s\eta(0, \cdot)} w(0, \cdot)\|_{L^2(\mathcal{O})}^2 = 2\text{Im} \left(\int_{(-T, 0) \times \mathcal{O}} M_1 \phi \bar{\phi} dt dx \right),$$

where M_1 stands for $M_{1,n}$.

Proof. In light of (3.24)-(3.25) it holds true that $\lim_{t \downarrow (-T)} \eta(t, x) = +\infty$ for all $x \in \mathcal{O}$, hence

$$\lim_{t \downarrow (-T)} \phi(t, x) = 0.$$

Therefore we have $J = \|\phi(0, \cdot)\|_{L^2(\mathcal{O})}^2 = \int_{(-T,0) \times \mathcal{O}} \partial_t |\phi|^2 dt dx$, from where we get that

$$J = 2\operatorname{Re} \left(\int_{(-T,0) \times \mathcal{O}} \partial_t \phi \bar{\phi} dt dx \right). \quad (4.46)$$

On the other hand, (3.26) and the Green formula yield

$$\begin{aligned} & \operatorname{Im} \left(\int_{(-T,0) \times \mathcal{O}} (M_1 \phi) \bar{\phi} dt dx \right) \\ &= \operatorname{Re} \left(\int_{(-T,0) \times \mathcal{O}} \partial_t \phi \bar{\phi} dt dx \right) + \operatorname{Im} \left(\int_{(-T,0) \times \mathcal{O}} \Delta \phi \bar{\phi} dt dx + s^2 \|\nabla \eta \phi\|_{L^2(-T,0) \times \mathcal{O}}^2 \right) \\ &= \operatorname{Re} \left(\int_{(-T,0) \times \mathcal{O}} \partial_t \phi \bar{\phi} dt dx \right) + \operatorname{Im} \left(\|\nabla \phi\|_{L^2((-T,0) \times \mathcal{O})}^2 \right) = \operatorname{Re} \left(\int_{(-T,0) \times \mathcal{O}} \partial_t \phi \bar{\phi} dt dx \right), \end{aligned}$$

so the result follows readily from this and (4.46). \square

Lemma 4.2. *Let w and J be the same as in Lemma 4.1. Then we have*

$$J \leq s^{-3/2} I(w), \quad s > 0,$$

where

$$I(w) := s \|e^{-s\eta} \nabla w\|_{L^2((-T,T) \times \mathcal{O})}^2 + s^3 \|e^{-s\eta} w\|_{L^2((-T,T) \times \mathcal{O})}^2 + \sum_{j=1,2} \|M_j e^{-s\eta} w\|_{L^2((-T,T) \times \mathcal{O})}^2, \quad (4.47)$$

the notation M_j , for $j = 1, 2$, being a shorthand for $M_{j,n}$.

Proof. In view of Lemma 4.1 and the Cauchy-Schwarz inequality, we have

$$\begin{aligned} J &\leq 2 \|M_1 \phi\|_{L^2((-T,T) \times \mathcal{O})} \|\phi\|_{L^2((-T,T) \times \mathcal{O})} \\ &\leq s^{-3/2} \left(\|M_1(e^{-s\eta} w)\|_{L^2((-T,T) \times \mathcal{O})}^2 + s^3 \|e^{-s\eta} w\|_{L^2((-T,T) \times \mathcal{O})}^2 \right). \end{aligned}$$

This and (4.47) yields the desired result. \square

4.1.2 Completion of the proof

The next step of the proof involves majorizing $I(w)$ with the aid of the Carleman inequality of Proposition 3.2. In view of (4.44)-(4.45) and (4.47) the estimate

$$\begin{aligned} I(w) &\leq C \left(s \|e^{-s\eta} \varphi^{1/2} (\partial_\nu \beta)^{1/2} \partial_\nu w\|_{L^2((-T, T) \times \tilde{\Gamma})}^2 + \|e^{-s\eta} (\chi f' - K v)\|_{L^2((-T, T) \times \mathcal{O})}^2 \right) \\ &\leq C \left(s \|e^{-s\eta} \varphi^{1/2} (\partial_\nu \beta)^{1/2} \partial_\nu w\|_{L^2((-T, T) \times \tilde{\Gamma})}^2 \right. \\ &\quad \left. + \sum_{j=0,1} \|e^{-s\eta} \nabla^j v\|_{L^2((-T, T) \times (\Omega_r \setminus \Omega_\ell))}^2 + \|e^{-s\eta} \chi f'\|_{L^2((-T, T) \times \Omega_r)}^2 \right), \end{aligned} \quad (4.48)$$

holds with $\tilde{\Gamma} := \Gamma_{\mathcal{O}}$ for $s \geq s_0$. Here and henceforth C denotes some generic positive constant. Bearing in mind that $\eta(0, x) = \inf_{t \in (-T, T)} \eta(t, x)$ for all $x \in \Omega$ and that $f' = (q_2 - q_1)u'_2$, we have

$$\|e^{-s\eta} \chi f'\|_{L^2((-T, T) \times \Omega_r)} \leq C \|e^{-s\eta(0, \cdot)} (q_1 - q_2)\|_{L^2(\Omega_\ell)}. \quad (4.49)$$

Here we used the fact that $\|u'_2\|_{L^\infty((-T, T) \times \Omega_\ell)} < \infty$. On the other hand, the function $\varphi \partial_\nu \beta$ being bounded in $(-T, T) \times \tilde{\Gamma}$ as well, we deduce from (4.48)-(4.49) that

$$\begin{aligned} I(w) &\leq C \left(s \|e^{-s\eta(0, \cdot)} \partial_\nu w\|_{L^2((-T, T) \times \tilde{\Gamma})}^2 + \sum_{j=0,1} \|e^{-s\eta(0, \cdot)} \nabla^j v\|_{L^2((-T, T) \times (\Omega_r \setminus \Omega_\ell))}^2 \right. \\ &\quad \left. + \|e^{-s\eta(0, \cdot)} (q_1 - q_2)\|_{L^2(\Omega_\ell)}^2 \right) \\ &\leq C \left(\mathbf{obs} + \|e^{-s\eta(0, \cdot)} (q_1 - q_2)\|_{L^2(\Omega_\ell)}^2 \right), \quad s \geq s_0, \end{aligned} \quad (4.50)$$

where we have set

$$\mathbf{obs} := s \|e^{-s\eta(0, \cdot)} \partial_\nu w\|_{L^2((-T, T) \times \tilde{\Gamma})}^2 + \sum_{j=0,1} \|e^{-s\eta(0, \cdot)} \nabla^j v\|_{L^2((-T, T) \times (\Omega_r \setminus \Omega_\ell))}^2. \quad (4.51)$$

In light of Lemma 4.2, (4.50)-(4.51) then yields

$$J = \|e^{-s\eta(0, \cdot)} \chi (q_1 - q_2) u_0\|_{L^2(\mathcal{O})}^2 \leq C s^{-3/2} \left(\mathbf{obs} + \|e^{-s\eta(0, \cdot)} (q_1 - q_2)\|_{L^2(\Omega_\ell)}^2 \right), \quad (4.52)$$

for all $s \geq s_0$. Further since $J \geq \alpha^2 \|e^{-s\eta(0, \cdot)} (q_1 - q_2)\|_{L^2(\Omega_\ell)}^2$ from the assumption (1.3) we get that

$$(\alpha^2 - C s^{-3/2}) \|e^{-s\eta(0, \cdot)} (q_1 - q_2)\|_{L^2(\Omega_\ell)}^2 \leq C s^{-3/2} \mathbf{obs}, \quad s \geq s_0, \quad (4.53)$$

from (4.52). Thus choosing $s \geq s_0$ so large that $\alpha^2 - C s^{-3/2} \geq \alpha^2/2$ and taking into account that $\inf_{x \in \Omega_\ell} e^{-s\eta(0, x)} > 0$, we derive from (4.51) and (4.53) that

$$\|q_1 - q_2\|_{L^2(\Omega_\ell)}^2 \leq C \left(\|\partial_\nu w\|_{L^2((-T, T) \times \tilde{\Gamma})}^2 + \sum_{j=0,1} \|\nabla^j v\|_{L^2((-T, T) \times (\Omega_r \setminus \Omega_\ell))}^2 \right). \quad (4.54)$$

Now, recalling from subsection 2.3 that $\|\nabla^j v\|_{L^2((-T,T)\times(\Omega_r\setminus\Omega_\ell))} = 2\|\nabla^j v\|_{L^2((0,T)\times(\Omega_r\setminus\Omega_\ell))}$ for $j = 0, 1$, and from the identity $w = \chi v$ that $\|\partial_\nu w\|_{L^2((-T,T)\times\tilde{\Gamma})} = 2\|\partial_\nu w\|_{L^2((0,T)\times\tilde{\Gamma})}$, we derive from (4.54) that

$$\|q_1 - q_2\|_{L^2(\Omega)}^2 \leq C \left(\|\partial_\nu w\|_{L^2((0,T)\times\tilde{\Gamma})}^2 + \sum_{j=0,1} \|\nabla^j v\|_{L^2((0,T)\times(\Omega_r\setminus\Omega_\ell))}^2 \right). \quad (4.55)$$

Here we used the identity $\|q_1 - q_2\|_{L^2(\Omega_\ell)} = \|q_1 - q_2\|_{L^2(\Omega)}$ arising from (1.4).

Last we set $\Gamma_* := \tilde{\Gamma} \cap (\partial\omega \times [-r, r])$ and $\Gamma_e := \tilde{\Gamma} \cap (\overline{\Omega}_L \setminus \overline{\Omega}_r)$ in such a way that

$$\tilde{\Gamma} = \Gamma_* \cup \Gamma_e, \quad \Gamma_* \cap \Gamma_e = \emptyset,$$

by (4.43). Since $\chi(x_n) = 0$ for every $|x_n| \geq r$ then the function $w = \chi v$ is necessarily uniformly zero in $(0, T) \times (\Omega_L \setminus \Omega_r)$. This entails

$$\partial_\nu w(t, \sigma) = 0, \quad (t, \sigma) \in (0, T) \times \Gamma_e.$$

As a consequence we have

$$\|\partial_\nu w\|_{L^2((0,T)\times\tilde{\Gamma})} = \|\partial_\nu w\|_{L^2((0,T)\times\Gamma_*)} \leq C \|\partial_\nu v\|_{L^2((0,T)\times\Gamma_*)}.$$

Putting this together with (4.55) and the identity $v = (u_1 - u_2)'$, we end up getting (1.7).

4.2 Proof of the stability estimate (1.8)

The proof of (1.8) is very similar to the one of (1.7). Therefore, in order to avoid the inadequate expense of the size of this article, we shall skip some technical details.

The function v still denoting the solution to (2.22), we put

$$I(v) := s \|e^{-s\eta} \nabla_{x'} v\|_{L^2(Q_e)}^2 + s^3 \|e^{-s\eta} v\|_{L^2(Q_e)}^2 + \sum_{j=1,2} \|M_j e^{-s\eta} v\|_{L^2(Q_e)}^2, \quad s > 0,$$

where we write M_j for $j = 1, 2$, instead of $M_{j,n}$. Then arguing in the exact same way as in the derivation of Lemmae 4.1-4.2, we get that

$$\|e^{-s\eta(0,\cdot)} v(0, \cdot)\|_{L^2(\Omega)}^2 = \|e^{-s\eta(0,\cdot)} (q_1 - q_2) u_0\|_{L^2(\Omega)}^2 \leq s^{-3/2} I(v), \quad (4.56)$$

for all $s > 0$.

On the other hand, applying Proposition 3.3 to (2.22) we obtain

$$I(v) \leq C \left(s \|e^{-s\eta} \varphi^{1/2} (\partial_\nu \beta)^{1/2} \partial_\nu v\|_{L^2((-T,T)\times\gamma_*\times\mathbb{R})}^2 + \|e^{-s\eta} f'\|_{L^2(Q_e)}^2 \right), \quad (4.57)$$

for every $s \geq s_0$. Further, taking into account that $\eta(0, x') = \inf_{t \in (-T, T)} \eta(t, x')$ for all $x' \in \omega$ and $f' = (q_2 - q_1)u'_2$, we derive from (1.6) that

$$\|e^{-s\eta} f'\|_{L^2(Q_e)} \leq C' \|e^{-s\eta(0, \cdot)}(q_1 - q_2)\|_{L^2(\Omega)}, \quad (4.58)$$

for some generic positive constant C' . Since $\varphi \partial_\nu \beta$ is bounded in $(-T, T) \times \gamma_* \times \mathbb{R}$, (4.57)-(4.58) then yield

$$I(v) \leq C' \left(s \|e^{-s\eta(0, \cdot)} \partial_\nu v\|_{L^2((-T, T) \times \gamma_* \times \mathbb{R})}^2 + \|e^{-s\eta(0, \cdot)}(q_1 - q_2)\|_{L^2(\Omega)}^2 \right), \quad s \geq s_0. \quad (4.59)$$

Putting (4.56) and (4.59) together we find that

$$\begin{aligned} & \|e^{-s\eta(0, \cdot)}(q_1 - q_2)u_0\|_{L^2(\Omega)}^2 \\ & \leq C' s^{-3/2} \left(s \|e^{-s\eta(0, \cdot)} \partial_\nu v\|_{L^2((-T, T) \times \gamma_* \times \mathbb{R})}^2 + \|e^{-s\eta(0, \cdot)}(q_1 - q_2)\|_{L^2(\Omega)}^2 \right), \quad s \geq s_0. \end{aligned}$$

From this and (1.3) then follows for any $s \geq s_0$ that

$$(\alpha^2 - C' s^{-3/2}) \|e^{-s\eta(0, \cdot)}(q_1 - q_2)\|_{L^2(\Omega)}^2 \leq C s^{-1/2} \|e^{-s\eta(0, \cdot)} \partial_\nu v\|_{L^2((-T, T) \times \gamma_* \times \mathbb{R})}^2. \quad (4.60)$$

Choosing $s \geq s_0$ so large that $\alpha^2 - C' s^{-3/2} \geq \alpha^2/2$ and using that $\inf_{x \in \omega} e^{-s\eta(0, x)} > 0$, we deduce from (4.60) that

$$\|q_1 - q_2\|_{L^2(\Omega)}^2 \leq C' \|\partial_\nu v\|_{L^2((-T, T) \times \gamma_* \times \mathbb{R})}^2.$$

Now (1.8) follows readily from this upon recalling from subsection 2.2 that $v = (u_1 - u_2)'$ and from subsection 2.3 that $\|\partial_\nu v\|_{L^2((-T, T) \times \gamma_* \times \mathbb{R})} = 2\|\partial_\nu v\|_{L^2((0, T) \times \gamma_* \times \mathbb{R})}$.

Acknowledgments. The authors are grateful to the unknown reviewers of this text for their useful suggestions and comments.

References

- [1] P. ALBANO, *Carleman estimates for the Euler-Bernoulli plate operator*, Elec. J. Diff. Equ. **53** (2000), 1-13.
- [2] C. BARDOS, G. LEBEAU, J. RAUCH, *Sharp sufficient conditions for the observation, control and stabilization from the boundary*, SIAM J. Control Optim. **30** (1992), 1024-1065.
- [3] L. BAUDOIN, J.-P. PUEL, *Uniqueness and stability in an inverse problem for the Schrödinger equation*, Inverse Probl., **18** (2002), 1537-1554.
- [4] M. BELLASSOUED, *Uniqueness and stability in determining the speed of propagation of second-order hyperbolic equation with variable coefficients*, Appl. Anal. **83** (2004), 983-1014.

- [5] M. BELLASSOUED, M. CHOULLI, *Logarithmic stability in the dynamical inverse problem for the Schrödinger equation by arbitrary boundary observation*, J. Math. Pures Appl. **91**(3) (2009), 233-255.
- [6] M. BELLASSOUED, M. CRISTOFOL, E. SOCCORSI, *Inverse boundary value problem for the dynamical heterogeneous Maxwell system*, Inverse Probl. **28**(9)(2012), 095009-26.
- [7] M. BELLASSOUED, M. YAMAMOTO, *Lipschitz stability in determining density and two Lamé coefficients*, J. Math. Anal. Appl. **329** (2007), 1240-1259.
- [8] A.L. BUKHGEIM, M.V. KLIBANOV, *Uniqueness in the large of a class of multi-dimensional inverse problems*, Sov. Math. Dokl. **17** (1981), 244-247.
- [9] A.L. BUKHGEIM, G. UHLMANN, *Recovering a potential from partial Cauchy data*, Commun. Part. Diff. Equat. **27** (2002), 653-668.
- [10] L. CARDOULIS, M. CRISTOFOL, P. GAITAN, *Inverse problems for the Schrödinger operator in an unbounded strip*, J. Inverse Ill-Posed Probl. **16**(2) (2008), 127-146.
- [11] P.-Y. CHANG, H.-H. LIN, *Conductance through a single impurity in the metallic zigzag carbon nanotube*, Appl. Phys. Lett. **95** (2009), 082104.
- [12] M. CHOULLI, E. SOCCORSI, *Recovering the twisting function in a twisted waveguide from the DN map*, arXiv:1209.5662.
- [13] M. CHOULLI, Y. KIAN, E. SOCCORSI, *Stable determination of time dependent scalar potential from boundary measurements in a periodic quantum waveguide*, arXiv:1306.6601.
- [14] M. CRISTOFOL, E. SOCCORSI, *Stability estimate in an inverse problem for non-autonomous magnetic Schrödinger equations*, Appl. Anal. **90**(10) (2011), 1499-1520.
- [15] O. YU. IMANUVILOV, *On Carleman estimates for hyperbolic equations*, Asympt. Anal. **32** (2002), 185-220.
- [16] O. YU. IMANUVILOV, M. YAMAMOTO, *Lipschitz stability in inverse parabolic problems by Carleman estimates*, Inverse Probl. **14** (1998), 1229-1249.
- [17] O. YU. IMANUVILOV, M. YAMAMOTO, *Global uniqueness and stability in determining coefficients of wave equations*, Commun. Part. Diff. Equat. **26** (2001), 1409-1425.
- [18] V. ISAKOV, *Uniqueness and stability in multi-dimensional inverse problems*, Inv. Prob. **9**(6) (1993),
- [19] V. ISAKOV, M. YAMAMOTO, *Carleman estimates with the Neumann boundary condition and its application to the observability inequality and inverse problems*, Contemp. Math. **268**(2000), 191-225.
- [20] C. KANE, L. BALENTS, M. P. A. FISHER, *Coulomb Interactions and Mesoscopic Effects in Carbon Nanotubes*, Phys. Rev. Lett. **79**(1997), 5086-5089.

- [21] C. E. KENIG, J. SJÖSTRAND, G. UHLMANN, *The Calderon problem with partial data*, Ann. Math. **165**(2007), 567-591.
- [22] Y. KIAN, Q. S. PHAN, E. SOCCORSI, *Hölder stable determination of a quantum scalar potential in unbounded cylindrical domains*, arXiv:1311.5323.
- [23] M. V. KLIBANOV, M. YAMAMOTO, *Lipschitz stability of an inverse problem for an acoustic equation*, Appl. Anal. **85** (2006), 515-538.
- [24] M. V. KLIBANOV, *Carleman estimates for global uniqueness, stability and numerical methods for inverse problems*, J. Inverse and Ill-Posed Problems **21** (2013), 477-560.
- [25] X. LI, G. UHLMANN, *Inverse problems with partial data in a slab*, Inverse Probl. and Imaging **4**(3) (2010), 449-462.
- [26] A. MERCADO, A. OSSES, L. ROSIER, *Inverse problems for the Schrödinger equation via Carleman inequalities with degenerate weights*, Inverse Probl. **24**(1) (2008), 015017.
- [27] D. TATARU, *Carleman estimates, unique continuation and controllability for anisotropic PDEs*, Contemp. Math. **209**(1) (1997), 267-279.
- [28] M. YAMAMOTO, *Uniqueness and stability in multidimensional hyperbolic inverse problems*, J. Math. Pures Appl. **78** (1) (1999), 65-98.
- [29] G. YUAN, M. YAMAMOTO, *Carleman estimates for the Schrödinger equation and applications to an inverse problem and an observability inequality*, Chin. Ann. Math. Ser. B **31** (2010), 555-578.